

# CHAPTER 5

## Integration

### EXERCISE SET 5.1

1. Endpoints  $0, \frac{1}{n}, \frac{2}{n}, \dots, \frac{n-1}{n}, 1$ ; using right endpoints

$$A_n = \left[ \sqrt{\frac{1}{n}} + \sqrt{\frac{2}{n}} + \dots + \sqrt{\frac{n-1}{n}} + 1 \right] \frac{1}{n}$$

$n$	2	5	10	50	100
$A_n$	0.853553	0.749739	0.710509	0.676095	0.671463

2. Endpoints  $0, \frac{1}{n}, \frac{2}{n}, \dots, \frac{n-1}{n}, 1$ ; using right endpoints

$$A_n = \left[ \frac{n}{n+1} + \frac{n}{n+2} + \frac{n}{n+3} + \dots + \frac{n}{2n-1} + \frac{1}{2} \right] \frac{1}{n}$$

$n$	2	5	10	50	100
$A_n$	0.583333	0.645635	0.668771	0.688172	0.690653

3. Endpoints  $0, \frac{\pi}{n}, \frac{2\pi}{n}, \dots, \frac{(n-1)\pi}{n}, \pi$ ; using right endpoints

$$A_n = [\sin(\pi/n) + \sin(2\pi/n) + \dots + \sin((n-1)\pi/n) + \sin \pi] \frac{\pi}{n}$$

$n$	2	5	10	50	100
$A_n$	1.57080	1.93376	1.98352	1.99935	1.99984

4. Endpoints  $0, \frac{\pi}{2n}, \frac{2\pi}{2n}, \dots, \frac{(n-1)\pi}{2n}, \frac{\pi}{2}$ ; using right endpoints

$$A_n = [\cos(\pi/2n) + \cos(2\pi/2n) + \dots + \cos((n-1)\pi/2n) + \cos(\pi/2)] \frac{\pi}{2n}$$

$n$	2	5	10	50	100
$A_n$	0.555359	0.834683	0.919405	0.984204	0.992120

5. Endpoints  $1, \frac{n+1}{n}, \frac{n+2}{n}, \dots, \frac{2n-1}{n}, 2$ ; using right endpoints

$$A_n = \left[ \frac{n}{n+1} + \frac{n}{n+2} + \dots + \frac{n}{2n-1} + \frac{1}{2} \right] \frac{1}{n}$$

$n$	2	5	10	50	100
$A_n$	0.583333	0.645635	0.668771	0.688172	0.690653

6. Endpoints  $-\frac{\pi}{2}, -\frac{\pi}{2} + \frac{\pi}{n}, -\frac{\pi}{2} + \frac{2\pi}{n}, \dots, -\frac{\pi}{2} + \frac{(n-1)\pi}{n}, \frac{\pi}{2}$ ; using right endpoints

$$A_n = \left[ \cos\left(-\frac{\pi}{2} + \frac{\pi}{n}\right) + \cos\left(-\frac{\pi}{2} + \frac{2\pi}{n}\right) + \dots + \cos\left(-\frac{\pi}{2} + \frac{(n-1)\pi}{n}\right) + \cos\left(\frac{\pi}{2}\right) \right] \frac{\pi}{n}$$

$n$	2	5	10	50	100
$A_n$	1.99985	1.93376	1.98352	1.99936	1.99985

7. Endpoints  $0, \frac{1}{n}, \frac{2}{n}, \dots, \frac{n-1}{n}, 1$ ; using right endpoints

$$A_n = \left[ \sqrt{1 - \left(\frac{1}{n}\right)^2} + \sqrt{1 - \left(\frac{2}{n}\right)^2} + \dots + \sqrt{1 - \left(\frac{n-1}{n}\right)^2} + 0 \right] \frac{1}{n}$$

$n$	2	5	10	50	100
$A_n$	0.433013	0.659262	0.726130	0.774567	0.780106

8. Endpoints  $-1, -1 + \frac{2}{n}, -1 + \frac{4}{n}, \dots, -1 + \frac{2(n-1)}{n}, 1$ ; using right endpoints

$$A_n = \left[ \sqrt{1 - \left(\frac{n-2}{n}\right)^2} + \sqrt{1 - \left(\frac{n-4}{n}\right)^2} + \dots + \sqrt{1 - \left(\frac{n-2}{n}\right)^2} + 0 \right] \frac{2}{n}$$

$n$	2	5	10	50	100
$A_n$	1	1.423837	1.518524	1.566097	1.569136

9.  $3(x-1)$

10.  $5(x-2)$

11.  $x(x+2)$

12.  $\frac{3}{2}(x-1)^2$

13.  $(x+3)(x-1)$

14.  $\frac{3}{2}x(x-2)$

15. The area in Exercise 13 is always 3 less than the area in Exercise 11. The regions are identical except that the area in Exercise 11 has the extra trapezoid with vertices at  $(0,0), (1,0), (0,2), (1,4)$  (with area 3).

16. (a) The region in question is a trapezoid, and the area of a trapezoid is  $\frac{1}{2}(h_1 + h_2)w$ .

$$\begin{aligned} \text{(b) From Part (a), } A'(x) &= \frac{1}{2}[f(a) + f(x)] + (x-a)\frac{1}{2}f'(x) \\ &= \frac{1}{2}[f(a) + f(x)] + (x-a)\frac{1}{2}\frac{f(x) - f(a)}{x-a} = f(x) \end{aligned}$$

17.  $B$  is also the area between the graph of  $f(x) = \sqrt{x}$  and the interval  $[0, 1]$  on the  $y$ -axis, so  $A + B$  is the area of the square.

18. If the plane is rotated about the line  $y = x$  then  $A$  becomes  $B$  and vice versa.

## EXERCISE SET 5.2

1. (a)  $\int \frac{x}{\sqrt{1+x^2}} dx = \sqrt{1+x^2} + C$       (b)  $\int x^2 \cos(1+x^3) dx = \frac{1}{3} \sin(1+x^3) + C$

2. (a)  $\frac{d}{dx}(\sin x - x \cos x + C) = \cos x - \cos x + x \sin x = x \sin x$

(b)  $\frac{d}{dx} \left( \frac{x}{\sqrt{1-x^2}} + C \right) = \frac{\sqrt{1-x^2} + x^2/\sqrt{1-x^2}}{1-x^2} = \frac{1}{(1-x^2)^{3/2}}$

3.  $\frac{d}{dx} \left[ \sqrt{x^3+5} \right] = \frac{3x^2}{2\sqrt{x^3+5}}$  so  $\int \frac{3x^2}{2\sqrt{x^3+5}} dx = \sqrt{x^3+5} + C$

4.  $\frac{d}{dx} \left[ \frac{x}{x^2 + 3} \right] = \frac{3 - x^2}{(x^2 + 3)^2}$  so  $\int \frac{3 - x^2}{(x^2 + 3)^2} dx = \frac{x}{x^2 + 3} + C$

5.  $\frac{d}{dx} [\sin(2\sqrt{x})] = \frac{\cos(2\sqrt{x})}{\sqrt{x}}$  so  $\int \frac{\cos(2\sqrt{x})}{\sqrt{x}} dx = \sin(2\sqrt{x}) + C$

6.  $\frac{d}{dx} [\sin x - x \cos x] = x \sin x$  so  $\int x \sin x dx = \sin x - x \cos x + C$

7. (a)  $x^9/9 + C$  (b)  $\frac{7}{12}x^{12/7} + C$  (c)  $\frac{2}{9}x^{9/2} + C$

8. (a)  $\frac{3}{5}x^{5/3} + C$  (b)  $-\frac{1}{5}x^{-5} + C = -\frac{1}{5x^5} + C$  (c)  $8x^{1/8} + C$

9. (a)  $\frac{1}{2} \int x^{-3} dx = -\frac{1}{4}x^{-2} + C$  (b)  $u^4/4 - u^2 + 7u + C$

10.  $\frac{3}{5}x^{5/3} - 5x^{4/5} + 4x + C$

11.  $\int (x^{-3} + x^{1/2} - 3x^{1/4} + x^2) dx = -\frac{1}{2}x^{-2} + \frac{2}{3}x^{3/2} - \frac{12}{5}x^{5/4} + \frac{1}{3}x^3 + C$

12.  $\int (7y^{-3/4} - y^{1/3} + 4y^{1/2}) dy = 28y^{1/4} - \frac{3}{4}y^{4/3} + \frac{8}{3}y^{3/2} + C$

13.  $\int (x + x^4) dx = x^2/2 + x^5/5 + C$

14.  $\int (4 + 4y^2 + y^4) dy = 4y + \frac{4}{3}y^3 + \frac{1}{5}y^5 + C$

15.  $\int x^{1/3}(4 - 4x + x^2) dx = \int (4x^{1/3} - 4x^{4/3} + x^{7/3}) dx = 3x^{4/3} - \frac{12}{7}x^{7/3} + \frac{3}{10}x^{10/3} + C$

16.  $\int (2 - x + 2x^2 - x^3) dx = 2x - \frac{1}{2}x^2 + \frac{2}{3}x^3 - \frac{1}{4}x^4 + C$

17.  $\int (x + 2x^{-2} - x^{-4}) dx = x^2/2 - 2/x + 1/(3x^3) + C$

18.  $\int (t^{-3} - 2) dt = -\frac{1}{2}t^{-2} - 2t + C$

19.  $-4 \cos x + 2 \sin x + C$  20.  $4 \tan x - \csc x + C$

21.  $\int (\sec^2 x + \sec x \tan x) dx = \tan x + \sec x + C$

22.  $\int (\sec x \tan x + 1) dx = \sec x + x + C$  23.  $\int \frac{\sec \theta}{\cos \theta} d\theta = \int \sec^2 \theta d\theta = \tan \theta + C$

24.  $\int \sin y dy = -\cos y + C$  25.  $\int \sec x \tan x dx = \sec x + C$



26.  $\int (\phi + 2 \csc^2 \phi) d\phi = \phi^2/2 - 2 \cot \phi + C$

27.  $\int (1 + \sin \theta) d\theta = \theta - \cos \theta + C$



28.  $\int \frac{2 \sin x \cos x}{\cos x} dx = 2 \int \sin x dx = -2 \cos x + C$

29.  $\int \frac{1 - \sin x}{1 - \sin^2 x} dx = \int \frac{1 - \sin x}{\cos^2 x} dx = \int (\sec^2 x - \sec x \tan x) dx = \tan x - \sec x + C$



30.  $\int \frac{1}{1 + \cos 2x} dx = \int \frac{1}{2 \cos^2 x} dx = \int \frac{1}{2} \sec^2 x dx = \frac{1}{2} \tan x + C$



(c)  $f(x) = x^2/2 - 1$



(c)  $y = \frac{2}{3}x^{3/2} - 2$



35.  $f'(x) = m = -\sin x$  so  $f(x) = \int (-\sin x) dx = \cos x + C$ ;  $f(0) = 2 = 1 + C$   
so  $C = 1$ ,  $f(x) = \cos x + 1$

36.  $f'(x) = m = (x+1)^2$ , so  $f(x) = \int (x+1)^2 dx = \frac{1}{3}(x+1)^3 + C$ ;

$$f(-2) = 8 = \frac{1}{3}(-2+1)^3 + C = -\frac{1}{3} + C, \quad 8 + \frac{1}{3} = \frac{25}{3}, \quad f(x) = \frac{1}{3}(x+1)^3 + \frac{25}{3}$$

37. (a)  $y(x) = \int x^{1/3} dx = \frac{3}{4}x^{4/3} + C$ ,  $y(1) = \frac{3}{4} + C = 2$ ,  $C = \frac{5}{4}$ ;  $y(x) = \frac{3}{4}x^{4/3} + \frac{5}{4}$

(b)  $y(t) = \int (\sin t + 1) dt = -\cos t + t + C$ ,  $y\left(\frac{\pi}{3}\right) = -\frac{1}{2} + \frac{\pi}{3} + C = \frac{1}{2}$ ,  $C = 1 - \frac{\pi}{3}$ ;  
 $y(t) = -\cos t + t + 1 - \frac{\pi}{3}$

(c)  $y(x) = \int (x^{1/2} + x^{-1/2}) dx = \frac{2}{3}x^{3/2} + 2x^{1/2} + C, y(1) = 0 = \frac{8}{3} + C, C = -\frac{8}{3},$   
 $y(x) = \frac{2}{3}x^{3/2} + 2x^{1/2} - \frac{8}{3}$

38. (a)  $y(x) = \int \frac{1}{8}x^{-3} dx = -\frac{1}{16}x^{-2} + C, y(1) = 0 = -\frac{1}{16} + C, C = \frac{1}{16}; y(x) = -\frac{1}{16}x^{-2} + \frac{1}{16}$   
(b)  $y(t) = \int (\sec^2 t - \sin t) dt = \tan t + \cos t + C, y(\frac{\pi}{4}) = 1 = 1 + \frac{\sqrt{2}}{2} + C, C = -\frac{\sqrt{2}}{2};$   
 $y(t) = \tan t + \cos t - \frac{\sqrt{2}}{2}$

(c)  $y(x) = \int x^{7/2} dx = \frac{2}{9}x^{9/2} + C, y(0) = 0 = C, C = 0; y(x) = \frac{2}{9}x^{9/2}$

39.  $f'(x) = \frac{2}{3}x^{3/2} + C_1; f(x) = \frac{4}{15}x^{5/2} + C_1x + C_2$

40.  $f'(x) = x^2/2 + \sin x + C_1$ , use  $f'(0) = 2$  to get  $C_1 = 2$  so  $f'(x) = x^2/2 + \sin x + 2$ ,  
 $f(x) = x^3/6 - \cos x + 2x + C_2$ , use  $f(0) = 1$  to get  $C_2 = 2$  so  $f(x) = x^3/6 - \cos x + 2x + 2$

41.  $dy/dx = 2x + 1, y = \int (2x + 1) dx = x^2 + x + C; y = 0$  when  $x = -3$   
so  $(-3)^2 + (-3) + C = 0, C = -6$  thus  $y = x^2 + x - 6$

42.  $dy/dx = x^2, y = \int x^2 dx = x^3/3 + C; y = 2$  when  $x = -1$  so  $(-1)^3/3 + C = 2, C = 7/3$   
thus  $y = x^3/3 + 7/3$

43.  $dy/dx = \int 6x dx = 3x^2 + C_1$ . The slope of the tangent line is  $-3$  so  $dy/dx = -3$  when  $x = 1$ .  
Thus  $3(1)^2 + C_1 = -3, C_1 = -6$  so  $dy/dx = 3x^2 - 6, y = \int (3x^2 - 6) dx = x^3 - 6x + C_2$ . If  $x = 1$ ,  
then  $y = 5 - 3(1) = 2$  so  $(1)^2 - 6(1) + C_2 = 2, C_2 = 7$  thus  $y = x^3 - 6x + 7$ .

44.  $dT/dx = C_1, T = C_1x + C_2; T = 25$  when  $x = 0$  so  $C_2 = 25, T = C_1x + 25$ .  $T = 85$  when  $x = 50$   
so  $50C_1 + 25 = 85, C_1 = 1.2, T = 1.2x + 25$

45. (a)  $F'(x) = G'(x) = 3x + 4$   
(b)  $F(0) = 16/6 = 8/3, G(0) = 0$ , so  $F(0) - G(0) = 8/3$   
(c)  $F(x) = (9x^2 + 24x + 16)/6 = 3x^2/2 + 4x + 8/3 = G(x) + 8/3$

46. (a)  $F'(x) = G'(x) = 10x/(x^2 + 5)^2$   
(b)  $F(0) = 0, G(0) = -1$ , so  $F(0) - G(0) = 1$   
(c)  $F(x) = \frac{x^2}{x^2 + 5} = \frac{(x^2 + 5) - 5}{x^2 + 5} = 1 - \frac{5}{x^2 + 5} = G(x) + 1$

47.  $\int (\sec^2 x - 1) dx = \tan x - x + C$                           48.  $\int (\csc^2 x - 1) dx = -\cot x - x + C$

49. (a)  $\frac{1}{2} \int (1 - \cos x) dx = \frac{1}{2}(x - \sin x) + C$                           (b)  $\frac{1}{2} \int (1 + \cos x) dx = \frac{1}{2}(x + \sin x) + C$

50. (a)  $F'(x) = G'(x) = f(x)$ , where  $f(x) = \begin{cases} 1, & x > 0 \\ -1, & x < 0 \end{cases}$

(b)  $G(x) - F(x) = \begin{cases} 2, & x > 0 \\ 3, & x < 0 \end{cases}$  so  $G(x) \neq F(x)$  plus a constant

(c) no, because  $(-\infty, 0) \cup (0, +\infty)$  is not an interval

51.  $v = \frac{1087}{2\sqrt{273}} \int T^{-1/2} dT = \frac{1087}{\sqrt{273}} T^{1/2} + C$ ,  $v(273) = 1087 = 1087 + C$  so  $C = 0$ ,  $v = \frac{1087}{\sqrt{273}} T^{1/2}$  ft/s

## EXERCISE SET 5.3

1. (a)  $\int u^{23} du = u^{24}/24 + C = (x^2 + 1)^{24}/24 + C$

(b)  $-\int u^3 du = -u^4/4 + C = -(\cos^4 x)/4 + C$

(c)  $2 \int \sin u du = -2 \cos u + C = -2 \cos \sqrt{x} + C$

(d)  $\frac{3}{8} \int u^{-1/2} du = \frac{3}{4} u^{1/2} + C = \frac{3}{4} \sqrt{4x^2 + 5} + C$

2. (a)  $\frac{1}{4} \int \sec^2 u du = \frac{1}{4} \tan u + C = \frac{1}{4} \tan(4x + 1) + C$

(b)  $\frac{1}{4} \int u^{1/2} du = \frac{1}{6} u^{3/2} + C = \frac{1}{6} (1 + 2y^2)^{3/2} + C$

(c)  $\frac{1}{\pi} \int u^{1/2} du = \frac{2}{3\pi} u^{3/2} + C = \frac{2}{3\pi} \sin^{3/2}(\pi\theta) + C$

(d)  $\int u^{4/5} du = \frac{5}{9} u^{9/5} + C = \frac{5}{9} (x^2 + 7x + 3)^{9/5} + C$

3. (a)  $-\int u du = -\frac{1}{2} u^2 + C = -\frac{1}{2} \cot^2 x + C$

(b)  $\int u^9 du = \frac{1}{10} u^{10} + C = \frac{1}{10} (1 + \sin t)^{10} + C$

(c)  $\frac{1}{2} \int \cos u du = \frac{1}{2} \sin u + C = \frac{1}{2} \sin 2x + C$

(d)  $\frac{1}{2} \int \sec^2 u du = \frac{1}{2} \tan u + C = \frac{1}{2} \tan x^2 + C$

4. (a)  $\int (u - 1)^2 u^{1/2} du = \int (u^{5/2} - 2u^{3/2} + u^{1/2}) du = \frac{2}{7} u^{7/2} - \frac{4}{5} u^{5/2} + \frac{2}{3} u^{3/2} + C$

$$= \frac{2}{7} (1 + x)^{7/2} - \frac{4}{5} (1 + x)^{5/2} + \frac{2}{3} (1 + x)^{3/2} + C$$

(b)  $\int \csc^2 u du = -\cot u + C = -\cot(\sin x) + C$

(c)  $\int \sin u du = -\cos u + C = -\cos(x - \pi) + C$

(d)  $\int \frac{du}{u^2} = -\frac{1}{u} + C = -\frac{1}{x^5 + 1} + C$

5.  $u = 2 - x^2$ ,  $du = -2x \, dx$ ;  $-\frac{1}{2} \int u^3 du = -u^4/8 + C = -(2 - x^2)^4/8 + C$

6.  $u = 3x - 1$ ,  $du = 3dx$ ;  $\frac{1}{3} \int u^5 du = \frac{1}{18}u^6 + C = \frac{1}{18}(3x - 1)^6 + C$

7.  $u = 8x$ ,  $du = 8dx$ ;  $\frac{1}{8} \int \cos u \, du = \frac{1}{8} \sin u + C = \frac{1}{8} \sin 8x + C$

8.  $u = 3x$ ,  $du = 3dx$ ;  $\frac{1}{3} \int \sin u \, du = -\frac{1}{3} \cos u + C = -\frac{1}{3} \cos 3x + C$

9.  $u = 4x$ ,  $du = 4dx$ ;  $\frac{1}{4} \int \sec u \tan u \, du = \frac{1}{4} \sec u + C = \frac{1}{4} \sec 4x + C$

10.  $u = 5x$ ,  $du = 5dx$ ;  $\frac{1}{5} \int \sec^2 u \, du = \frac{1}{5} \tan u + C = \frac{1}{5} \tan 5x + C$

11.  $u = 7t^2 + 12$ ,  $du = 14t \, dt$ ;  $\frac{1}{14} \int u^{1/2} du = \frac{1}{21}u^{3/2} + C = \frac{1}{21}(7t^2 + 12)^{3/2} + C$

12.  $u = 4 - 5x^2$ ,  $du = -10x \, dx$ ;  $-\frac{1}{10} \int u^{-1/2} du = -\frac{1}{5}u^{1/2} + C = -\frac{1}{5}\sqrt{4 - 5x^2} + C$

13.  $u = x^3 + 1$ ,  $du = 3x^2 dx$ ;  $\frac{1}{3} \int u^{-1/2} du = \frac{2}{3}u^{1/2} + C = \frac{2}{3}\sqrt{x^3 + 1} + C$

14.  $u = 1 - 3x$ ,  $du = -3dx$ ;  $-\frac{1}{3} \int u^{-2} du = \frac{1}{3}u^{-1} + C = \frac{1}{3}(1 - 3x)^{-1} + C$

15.  $u = 4x^2 + 1$ ,  $du = 8x \, dx$ ;  $\frac{1}{8} \int u^{-3} du = -\frac{1}{16}u^{-2} + C = -\frac{1}{16}(4x^2 + 1)^{-2} + C$

16.  $u = 3x^2$ ,  $du = 6x \, dx$ ;  $\frac{1}{6} \int \cos u \, du = \frac{1}{6} \sin u + C = \frac{1}{6} \sin(3x^2) + C$

17.  $u = 5/x$ ,  $du = -(5/x^2)dx$ ;  $-\frac{1}{5} \int \sin u \, du = \frac{1}{5} \cos u + C = \frac{1}{5} \cos(5/x) + C$

18.  $u = \sqrt{x}$ ,  $du = \frac{1}{2\sqrt{x}}dx$ ;  $2 \int \sec^2 u \, du = 2 \tan u + C = 2 \tan \sqrt{x} + C$

19.  $u = x^3$ ,  $du = 3x^2 dx$ ;  $\frac{1}{3} \int \sec^2 u \, du = \frac{1}{3} \tan u + C = \frac{1}{3} \tan(x^3) + C$

20.  $u = \cos 2t$ ,  $du = -2 \sin 2t \, dt$ ;  $-\frac{1}{2} \int u^3 du = -\frac{1}{8}u^4 + C = -\frac{1}{8} \cos^4 2t + C$

21.  $u = \sin 3t$ ,  $du = 3 \cos 3t \, dt$ ;  $\frac{1}{3} \int u^5 du = \frac{1}{18}u^6 + C = \frac{1}{18} \sin^6 3t + C$

22.  $u = 5 + \cos 2\theta$ ,  $du = -2 \sin 2\theta \, d\theta$ ;  $-\frac{1}{2} \int u^{-3} du = \frac{1}{4}u^{-2} + C = \frac{1}{4}(5 + \cos 2\theta)^{-2} + C$

**23.**  $u = 2 - \sin 4\theta$ ,  $du = -4 \cos 4\theta d\theta$ ;  $-\frac{1}{4} \int u^{1/2} du = -\frac{1}{6} u^{3/2} + C = -\frac{1}{6} (2 - \sin 4\theta)^{3/2} + C$

**24.**  $u = \tan 5x$ ,  $du = 5 \sec^2 5x dx$ ;  $\frac{1}{5} \int u^3 du = \frac{1}{20} u^4 + C = \frac{1}{20} \tan^4 5x + C$

**25.**  $u = \sec 2x$ ,  $du = 2 \sec 2x \tan 2x dx$ ;  $\frac{1}{2} \int u^2 du = \frac{1}{6} u^3 + C = \frac{1}{6} \sec^3 2x + C$

**26.**  $u = \sin \theta$ ,  $du = \cos \theta d\theta$ ;  $\int \sin u du = -\cos u + C = -\cos(\sin \theta) + C$

**27.**  $u = x - 3$ ,  $x = u + 3$ ,  $dx = du$

$$\int (u+3)u^{1/2} du = \int (u^{3/2} + 3u^{1/2}) du = \frac{2}{5} u^{5/2} + 2u^{3/2} + C = \frac{2}{5} (x-3)^{5/2} + 2(x-3)^{3/2} + C$$

**28.**  $u = y + 1$ ,  $y = u - 1$ ,  $dy = du$

$$\int \frac{u-1}{u^{1/2}} du = \int (u^{1/2} - u^{-1/2}) du = \frac{2}{3} u^{3/2} - 2u^{1/2} + C = \frac{2}{3} (y+1)^{3/2} - 2(y+1)^{1/2} + C$$

**29.**  $\int \sin^2 2\theta \sin 2\theta d\theta = \int (1 - \cos^2 2\theta) \sin 2\theta d\theta$ ;  $u = \cos 2\theta$ ,  $du = -2 \sin 2\theta d\theta$ ,

$$-\frac{1}{2} \int (1 - u^2) du = -\frac{1}{2} u + \frac{1}{6} u^3 + C = -\frac{1}{2} \cos 2\theta + \frac{1}{6} \cos^3 2\theta + C$$

**30.**  $\sec^2 3\theta = \tan^2 3\theta + 1$ ,  $u = 3\theta$ ,  $du = 3d\theta$

$$\int \sec^4 3\theta d\theta = \frac{1}{3} \int (\tan^2 u + 1) \sec^2 u du = \frac{1}{9} \tan^3 u + \frac{1}{3} \tan u + C = \frac{1}{9} \tan^3 3\theta + \frac{1}{3} \tan 3\theta + C$$

**31.**  $u = a + bx$ ,  $du = bdx$ ,

$$\int (a + bx)^n dx = \frac{1}{b} \int u^n du = \frac{(a + bx)^{n+1}}{b(n+1)} + C$$

**32.**  $u = a + bx$ ,  $du = b dx$ ,  $dx = \frac{1}{b} du$

$$\frac{1}{b} \int u^{1/n} du = \frac{n}{b(n+1)} u^{(n+1)/n} + C = \frac{n}{b(n+1)} (a + bx)^{(n+1)/n} + C$$

**33.**  $u = \sin(a + bx)$ ,  $du = b \cos(a + bx) dx$

$$\frac{1}{b} \int u^n du = \frac{1}{b(n+1)} u^{n+1} + C = \frac{1}{b(n+1)} \sin^{n+1}(a + bx) + C$$

**35. (a)** with  $u = \sin x$ ,  $du = \cos x dx$ ;  $\int u du = \frac{1}{2} u^2 + C_1 = \frac{1}{2} \sin^2 x + C_1$ ;

with  $u = \cos x$ ,  $du = -\sin x dx$ ;  $-\int u du = -\frac{1}{2} u^2 + C_2 = -\frac{1}{2} \cos^2 x + C_2$

**(b)** because they differ by a constant:

$$\left( \frac{1}{2} \sin^2 x + C_1 \right) - \left( -\frac{1}{2} \cos^2 x + C_2 \right) = \frac{1}{2} (\sin^2 x + \cos^2 x) + C_1 - C_2 = 1/2 + C_1 - C_2$$

36. (a) First method:  $\int (25x^2 - 10x + 1)dx = \frac{25}{3}x^3 - 5x^2 + x + C_1$ ;

second method:  $\frac{1}{5} \int u^2 du = \frac{1}{15}u^3 + C_2 = \frac{1}{15}(5x - 1)^3 + C_2$

(b)  $\frac{1}{15}(5x - 1)^3 + C_2 = \frac{1}{15}(125x^3 - 75x^2 + 15x - 1) + C_2 = \frac{25}{3}x^3 - 5x^2 + x - \frac{1}{15} + C_2$ ;

the answers differ by a constant.

37.  $y(x) = \int \sqrt{3x + 1} dx = \frac{2}{9}(3x + 1)^{3/2} + C$ ,

$$y(1) = \frac{16}{9} + C = 5, C = \frac{29}{9} \text{ so } y(x) = \frac{2}{9}(3x + 1)^{3/2} + \frac{29}{9}$$


38.  $y(x) = \int (6 - 5 \sin 2x) dx = 6x + \frac{5}{2} \cos 2x + C$ ,

$$y(0) = \frac{5}{2} + C = 3, C = \frac{1}{2} \text{ so } y(x) = 6x + \frac{5}{2} \cos 2x + \frac{1}{2}$$

39.  $f'(x) = m = \sqrt{3x + 1}, f(x) = \int (3x + 1)^{1/2} dx = \frac{2}{9}(3x + 1)^{3/2} + C; f(0) = 1 = \frac{2}{9} + C, C = \frac{7}{9}$ ,

$$\text{so } f(x) = \frac{2}{9}(3x + 1)^{3/2} + \frac{7}{9}$$

40.



41.  $p(t) = \int (4 + 0.15t)^{3/2} dt = \frac{8}{3}(4 + 0.15t)^{5/2} + C; p(0) = 100,000 = \frac{8}{3}4^{5/2} + C = \frac{256}{3} + C$ ,

$$C = 100,000 - \frac{256}{3} \approx 99,915, p(t) \approx \frac{8}{3}(4 + 0.15t)^{5/2} + 99,915, p(5) \approx \frac{8}{3}(4.75)^{5/2} + 99,915 \approx 100,046$$

## EXERCISE SET 5.4

1. (a)  $1 + 8 + 27 = 36$

(b)  $5 + 8 + 11 + 14 + 17 = 55$

(c)  $20 + 12 + 6 + 2 + 0 + 0 = 40$

(d)  $1 + 1 + 1 + 1 + 1 + 1 = 6$

(e)  $1 - 2 + 4 - 8 + 16 = 11$

(f)  $0 + 0 + 0 + 0 + 0 + 0 = 0$

2. (a)  $1 + 0 - 3 + 0 = -2$

(b)  $1 - 1 + 1 - 1 + 1 - 1 = 0$

(c)  $\pi^2 + \pi^2 + \dots + \pi^2 = 14\pi^2$   
(14 terms)

(d)  $2^4 + 2^5 + 2^6 = 112$

(e)  $\sqrt{1} + \sqrt{2} + \sqrt{3} + \sqrt{4} + \sqrt{5} + \sqrt{6}$

(f)  $1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 - 1 + 1 = 1$

3.  $\sum_{k=1}^{10} k$

4.  $\sum_{k=1}^{20} 3k$

5.  $\sum_{k=1}^{10} 2k$

6.  $\sum_{k=1}^8 (2k - 1)$

7.  $\sum_{k=1}^6 (-1)^{k+1} (2k - 1)$

8.  $\sum_{k=1}^5 (-1)^{k+1} \frac{1}{k}$

9. (a)  $\sum_{k=1}^{50} 12k$

(b)  $\sum_{k=1}^{50} (2k - 1)$

10. (a)  $\sum_{k=1}^5 (-1)^{k+1} a_k$       (b)  $\sum_{k=0}^5 (-1)^{k+1} b_k$       (c)  $\sum_{k=0}^n a_k x^k$       (d)  $\sum_{k=0}^5 a^{5-k} b^k$

11.  $\frac{1}{2}(100)(100 + 1) = 5050$

12.  $7 \sum_{k=1}^{100} k + \sum_{k=1}^{100} 1 = \frac{7}{2}(100)(101) + 100 = 35,450$

13.  $\frac{1}{6}(20)(21)(41) = 2870$

14.  $\sum_{k=1}^{20} k^2 - \sum_{k=1}^3 k^2 = 2870 - 14 = 2856$

15.  $\sum_{k=1}^{30} k(k^2 - 4) = \sum_{k=1}^{30} (k^3 - 4k) = \sum_{k=1}^{30} k^3 - 4 \sum_{k=1}^{30} k = \frac{1}{4}(30)^2(31)^2 - 4 \cdot \frac{1}{2}(30)(31) = 214,365$

16.  $\sum_{k=1}^6 k - \sum_{k=1}^6 k^3 = \frac{1}{2}(6)(7) - \frac{1}{4}(6)^2(7)^2 = -420$

17.  $\sum_{k=1}^n \frac{3k}{n} = \frac{3}{n} \sum_{k=1}^n k = \frac{3}{n} \cdot \frac{1}{2} n(n + 1) = \frac{3}{2}(n + 1)$

18.  $\sum_{k=1}^{n-1} \frac{k^2}{n} = \frac{1}{n} \sum_{k=1}^{n-1} k^2 = \frac{1}{n} \cdot \frac{1}{6}(n - 1)(n)(2n - 1) = \frac{1}{6}(n - 1)(2n - 1)$

19.  $\sum_{k=1}^{n-1} \frac{k^3}{n^2} = \frac{1}{n^2} \sum_{k=1}^{n-1} k^3 = \frac{1}{n^2} \cdot \frac{1}{4}(n - 1)^2 n^2 = \frac{1}{4}(n - 1)^2$

20.  $\sum_{k=1}^n \left( \frac{5}{n} - \frac{2k}{n} \right) = \frac{5}{n} \sum_{k=1}^n 1 - \frac{2}{n} \sum_{k=1}^n k = \frac{5}{n}(n) - \frac{2}{n} \cdot \frac{1}{2} n(n + 1) = 4 - n$

22.  $\frac{n(n + 1)}{2} = 465, n^2 + n - 930 = 0, (n + 31)(n - 30) = 0, n = 30.$

23.  $\frac{1 + 2 + 3 + \cdots + n}{n^2} = \sum_{k=1}^n \frac{k}{n^2} = \frac{1}{n^2} \sum_{k=1}^n k = \frac{1}{n^2} \cdot \frac{1}{2} n(n + 1) = \frac{n + 1}{2n}; \lim_{n \rightarrow +\infty} \frac{n + 1}{2n} = \frac{1}{2}$

24.  $\frac{1^2 + 2^2 + 3^2 + \cdots + n^2}{n^3} = \sum_{k=1}^n \frac{k^2}{n^3} = \frac{1}{n^3} \sum_{k=1}^n k^2 = \frac{1}{n^3} \cdot \frac{1}{6} n(n+1)(2n+1) = \frac{(n+1)(2n+1)}{6n^2};$

$$\lim_{n \rightarrow +\infty} \frac{(n+1)(2n+1)}{6n^2} = \lim_{n \rightarrow +\infty} \frac{1}{6}(1+1/n)(2+1/n) = \frac{1}{3}$$

25.  $\sum_{k=1}^n \frac{5k}{n^2} = \frac{5}{n^2} \sum_{k=1}^n k = \frac{5}{n^2} \cdot \frac{1}{2} n(n+1) = \frac{5(n+1)}{2n}; \lim_{n \rightarrow +\infty} \frac{5(n+1)}{2n} = \frac{5}{2}$

26.  $\sum_{k=1}^{n-1} \frac{2k^2}{n^3} = \frac{2}{n^3} \sum_{k=1}^{n-1} k^2 = \frac{2}{n^3} \cdot \frac{1}{6} (n-1)n(2n-1) = \frac{(n-1)(2n-1)}{3n^2};$

$$\lim_{n \rightarrow +\infty} \frac{(n-1)(2n-1)}{3n^2} = \lim_{n \rightarrow +\infty} \frac{1}{3}(1-1/n)(2-1/n) = \frac{2}{3}$$

27. (a)  $\sum_{j=0}^5 2^j$       (b)  $\sum_{j=1}^6 2^{j-1}$       (c)  $\sum_{j=2}^7 2^{j-2}$

28. (a)  $\sum_{k=1}^5 (k+4)2^{k+8}$       (b)  $\sum_{k=9}^{13} (k-4)2^k$

29. Endpoints 2, 3, 4, 5, 6;  $\Delta x = 1$ ;

(a) Left endpoints:  $\sum_{k=1}^4 f(x_k^*) \Delta x = 7 + 10 + 13 + 16 = 46$

(b) Midpoints:  $\sum_{k=1}^4 f(x_k^*) \Delta x = 8.5 + 11.5 + 14.5 + 17.5 = 52$

(c) Right endpoints:  $\sum_{k=1}^4 f(x_k^*) \Delta x = 10 + 13 + 16 + 19 = 58$

30. Endpoints 1, 3, 5, 7, 9,  $\Delta x = 2$ ;

(a) Left endpoints:  $\sum_{k=1}^4 f(x_k^*) \Delta x = \left(1 + \frac{1}{3} + \frac{1}{5} + \frac{1}{7}\right) 2 = \frac{352}{105}$

(b) Midpoints:  $\sum_{k=1}^4 f(x_k^*) \Delta x = \left(\frac{1}{2} + \frac{1}{4} + \frac{1}{6} + \frac{1}{8}\right) 2 = \frac{25}{12}$

(c) Right endpoints:  $\sum_{k=1}^4 f(x_k^*) \Delta x = \left(\frac{1}{3} + \frac{1}{5} + \frac{1}{7} + \frac{1}{9}\right) 2 = \frac{496}{315}$

31. Endpoints: 0,  $\pi/4$ ,  $\pi/2$ ,  $3\pi/4$ ,  $\pi$ ;  $\Delta x = \pi/4$

(a) Left endpoints:  $\sum_{k=1}^4 f(x_k^*) \Delta x = \left(1 + \sqrt{2}/2 + 0 - \sqrt{2}/2\right) (\pi/4) = \pi/4$

(b) Midpoints:  $\sum_{k=1}^4 f(x_k^*) \Delta x = [\cos(\pi/8) + \cos(3\pi/8) + \cos(5\pi/8) + \cos(7\pi/8)] (\pi/4)$   
 $= [\cos(\pi/8) + \cos(3\pi/8) - \cos(3\pi/8) - \cos(\pi/8)] (\pi/4) = 0$

(c) Right endpoints:  $\sum_{k=1}^4 f(x_k^*) \Delta x = \left(\sqrt{2}/2 + 0 - \sqrt{2}/2 - 1\right) (\pi/4) = -\pi/4$

**32.** Endpoints  $-1, 0, 1, 2, 3; \Delta x = 1$

(a)  $\sum_{k=1}^4 f(x_k^*) \Delta x = -3 + 0 + 1 + 0 = -2$

(b)  $\sum_{k=1}^4 f(x_k^*) \Delta x = -\frac{5}{4} + \frac{3}{4} + \frac{3}{4} + \frac{15}{4} = 4$

(c)  $\sum_{k=1}^4 f(x_k^*) \Delta x = 0 + 1 + 0 - 3 = -2$

**33.** (a) 0.718771403, 0.705803382, 0.698172179

(b) 0.668771403, 0.680803382, 0.688172179

(c) 0.692835360, 0.693069098, 0.693134682

**34.** (a) 0.761923639, 0.712712753, 0.684701150

(b) 0.584145862, 0.623823864, 0.649145594

(c) 0.663501867, 0.665867079, 0.666538346

**35.** (a) 4.884074734, 5.115572731, 5.248762738

(b) 5.684074734, 5.515572731, 5.408762738

(c) 5.34707029, 5.338362719, 5.334644416

**36.** (a) 0.919403170, 0.960215997, 0.984209789

(b) 1.076482803, 1.038755813, 1.015625715

(c) 1.001028824, 1.000257067, 1.000041125

**37.**  $\Delta x = \frac{3}{n}$ ,  $x_k^* = 1 + \frac{3}{n}k$ ;  $f(x_k^*) \Delta x = \frac{1}{2}x_k^* \Delta x = \frac{1}{2} \left(1 + \frac{3}{n}k\right) \frac{3}{n} = \frac{3}{2} \left[\frac{1}{n} + \frac{3}{n^2}k\right]$

$$\sum_{k=1}^n f(x_k^*) \Delta x = \frac{3}{2} \left[ \sum_{k=1}^n \frac{1}{n} + \sum_{k=1}^n \frac{3}{n^2}k \right] = \frac{3}{2} \left[ 1 + \frac{3}{n^2} \cdot \frac{1}{2}n(n+1) \right] = \frac{3}{2} \left[ 1 + \frac{3}{2} \frac{n+1}{n} \right]$$

$$A = \lim_{n \rightarrow +\infty} \frac{3}{2} \left[ 1 + \frac{3}{2} \left(1 + \frac{1}{n}\right) \right] = \frac{3}{2} \left(1 + \frac{3}{2}\right) = \frac{15}{4}$$

**38.**  $\Delta x = \frac{5}{n}$ ,  $x_k^* = 0 + k\frac{5}{n}$ ;  $f(x_k^*) \Delta x = (5 - x_k^*) \Delta x = \left(5 - \frac{5}{n}k\right) \frac{5}{n} = \frac{25}{n} - \frac{25}{n^2}k$

$$\sum_{k=1}^n f(x_k^*) \Delta x = \sum_{k=1}^n \frac{25}{n} - \frac{25}{n^2} \sum_{k=1}^n k = 25 - \frac{25}{n^2} \cdot \frac{1}{2}n(n+1) = 25 - \frac{25}{2} \left(\frac{n+1}{n}\right)$$

$$A = \lim_{n \rightarrow +\infty} \left[ 25 - \frac{25}{2} \left(1 + \frac{1}{n}\right) \right] = 25 - \frac{25}{2} = \frac{25}{2}$$

**39.**  $\Delta x = \frac{3}{n}$ ,  $x_k^* = 0 + k\frac{3}{n}$ ;  $f(x_k^*) \Delta x = \left(9 - 9\frac{k^2}{n^2}\right) \frac{3}{n}$

$$\sum_{k=1}^n f(x_k^*) \Delta x = \sum_{k=1}^n \left(9 - 9\frac{k^2}{n^2}\right) \frac{3}{n} = \frac{27}{n} \sum_{k=1}^n \left(1 - \frac{k^2}{n^2}\right) = 27 - \frac{27}{n^3} \sum_{k=1}^n k^2$$

$$A = \lim_{n \rightarrow +\infty} \left[ 27 - \frac{27}{n^3} \sum_{k=1}^n k^2 \right] = 27 - 27 \left(\frac{1}{3}\right) = 18$$

40.  $\Delta x = \frac{3}{n}$ ,  $x_k^* = k \frac{3}{n}$

$$\begin{aligned} f(x_k^*)\Delta x &= \left[4 - \frac{1}{4}(x_k^*)^2\right]\Delta x = \left[4 - \frac{1}{4}\frac{9k^2}{n^2}\right]\frac{3}{n} = \frac{12}{n} - \frac{27k^2}{4n^3} \\ \sum_{k=1}^n f(x_k^*)\Delta x &= \sum_{k=1}^n \frac{12}{n} - \frac{27}{4n^3} \sum_{k=1}^n k^2 \\ &= 12 - \frac{27}{4n^3} \cdot \frac{1}{6}n(n+1)(2n+1) = 12 - \frac{9}{8} \frac{(n+1)(2n+1)}{n^2} \\ A &= \lim_{n \rightarrow +\infty} \left[12 - \frac{9}{8} \left(1 + \frac{1}{n}\right) \left(2 + \frac{1}{n}\right)\right] = 12 - \frac{9}{8}(1)(2) = 39/4 \end{aligned}$$

41.  $\Delta x = \frac{4}{n}$ ,  $x_k^* = 2 + k \frac{4}{n}$

$$\begin{aligned} f(x_k^*)\Delta x &= (x_k^*)^3 \Delta x = \left[2 + \frac{4}{n}k\right]^3 \frac{4}{n} = \frac{32}{n} \left[1 + \frac{2}{n}k\right]^3 = \frac{32}{n} \left[1 + \frac{6}{n}k + \frac{12}{n^2}k^2 + \frac{8}{n^3}k^3\right] \\ \sum_{k=1}^n f(x_k^*)\Delta x &= \frac{32}{n} \left[\sum_{k=1}^n 1 + \frac{6}{n} \sum_{k=1}^n k + \frac{12}{n^2} \sum_{k=1}^n k^2 + \frac{8}{n^3} \sum_{k=1}^n k^3\right] \\ &= \frac{32}{n} \left[n + \frac{6}{n} \cdot \frac{1}{2}n(n+1) + \frac{12}{n^2} \cdot \frac{1}{6}n(n+1)(2n+1) + \frac{8}{n^3} \cdot \frac{1}{4}n^2(n+1)^2\right] \\ &= 32 \left[1 + 3\frac{n+1}{n} + 2\frac{(n+1)(2n+1)}{n^2} + 2\frac{(n+1)^2}{n^2}\right] \\ A &= \lim_{n \rightarrow +\infty} 32 \left[1 + 3\left(1 + \frac{1}{n}\right) + 2\left(1 + \frac{1}{n}\right)\left(2 + \frac{1}{n}\right) + 2\left(1 + \frac{1}{n}\right)^2\right] \\ &= 32[1 + 3(1) + 2(1)(2) + 2(1)^2] = 320 \end{aligned}$$

42.  $\Delta x = \frac{2}{n}$ ,  $x_k^* = -3 + k \frac{2}{n}$ ;  $f(x_k^*)\Delta x = [1 - (x_k^*)^3]\Delta x = \left[1 - \left(-3 + \frac{2}{n}k\right)^3\right] \frac{2}{n}$

$$\begin{aligned} \sum_{k=1}^n f(x_k^*)\Delta x &= \frac{2}{n} \left[28n - 27(n+1) + 6\frac{(n+1)(2n+1)}{n} - 2\frac{(n+1)^2}{n}\right] \\ A &= \lim_{n \rightarrow +\infty} 2 \left[28 - 27\left(1 + \frac{1}{n}\right) + 6\left(1 + \frac{1}{n}\right)\left(2 + \frac{1}{n}\right) - 2\left(1 + \frac{1}{n}\right)^2\right] \\ &= 2(28 - 27 + 12 - 2) = 22 \end{aligned}$$

43.  $\Delta x = \frac{3}{n}$ ,  $x_k^* = 1 + (k-1)\frac{3}{n}$

$$\begin{aligned} f(x_k^*)\Delta x &= \frac{1}{2}x_k^*\Delta x = \frac{1}{2} \left[1 + (k-1)\frac{3}{n}\right] \frac{3}{n} = \frac{1}{2} \left[\frac{3}{n} + (k-1)\frac{9}{n^2}\right] \\ \sum_{k=1}^n f(x_k^*)\Delta x &= \frac{1}{2} \left[\sum_{k=1}^n \frac{3}{n} + \frac{9}{n^2} \sum_{k=1}^n (k-1)\right] = \frac{1}{2} \left[3 + \frac{9}{n^2} \cdot \frac{1}{2}(n-1)n\right] = \frac{3}{2} + \frac{9}{4} \frac{n-1}{n} \\ A &= \lim_{n \rightarrow +\infty} \left[\frac{3}{2} + \frac{9}{4} \left(1 - \frac{1}{n}\right)\right] = \frac{3}{2} + \frac{9}{4} = \frac{15}{4} \end{aligned}$$

44.  $\Delta x = \frac{5}{n}$ ,  $x_k^* = \frac{5}{n}(k - 1)$

$$f(x_k^*)\Delta x = (5 - x_k^*)\Delta x = \left[5 - \frac{5}{n}(k - 1)\right] \frac{5}{n} = \frac{25}{n} - \frac{25}{n^2}(k - 1)$$

$$\sum_{k=1}^n f(x_k^*)\Delta x = \frac{25}{n} \sum_{k=1}^n 1 - \frac{25}{n^2} \sum_{k=1}^n (k - 1) = 25 - \frac{25}{2} \frac{n-1}{n}$$

$$A = \lim_{n \rightarrow +\infty} \left[ 25 - \frac{25}{2} \left(1 - \frac{1}{n}\right) \right] = 25 - \frac{25}{2} = \frac{25}{2}$$

45.  $\Delta x = \frac{3}{n}$ ,  $x_k^* = 0 + (k - 1)\frac{3}{n}$ ;  $f(x_k^*)\Delta x = (9 - 9\frac{(k-1)^2}{n^2})\frac{3}{n}$

$$\sum_{k=1}^n f(x_k^*)\Delta x = \sum_{k=1}^n \left[ 9 - 9\frac{(k-1)^2}{n^2} \right] \frac{3}{n} = \frac{27}{n} \sum_{k=1}^n \left( 1 - \frac{(k-1)^2}{n^2} \right) = 27 - \frac{27}{n^3} \sum_{k=1}^n k^2 + \frac{54}{n^3} \sum_{k=1}^n k - \frac{27}{n^2}$$

$$A = \lim_{n \rightarrow +\infty} = 27 - 27 \left(\frac{1}{3}\right) + 0 + 0 = 18$$

46.  $\Delta x = \frac{3}{n}$ ,  $x_k^* = (k - 1)\frac{3}{n}$

$$f(x_k^*)\Delta x = \left[ 4 - \frac{1}{4}(x_k^*)^2 \right] \Delta x = \left[ 4 - \frac{1}{4} \frac{9(k-1)^2}{n^2} \right] \frac{3}{n} = \frac{12}{n} - \frac{27k^2}{4n^3} + \frac{27k}{2n^3} - \frac{27}{4n^3}$$

$$\begin{aligned} \sum_{k=1}^n f(x_k^*)\Delta x &= \sum_{k=1}^n \frac{12}{n} - \frac{27}{4n^3} \sum_{k=1}^n k^2 + \frac{27}{2n^3} \sum_{k=1}^n k - \frac{27}{4n^3} \sum_{k=1}^n 1 \\ &= 12 - \frac{27}{4n^3} \cdot \frac{1}{6} n(n+1)(2n+1) + \frac{27}{2n^3} \frac{n(n+1)}{2} - \frac{27}{4n^2} \\ &= 12 - \frac{9}{8} \frac{(n+1)(2n+1)}{n^2} + \frac{27}{4n} + \frac{27}{4n^2} - \frac{27}{4n^2} \end{aligned}$$

$$A = \lim_{n \rightarrow +\infty} \left[ 12 - \frac{9}{8} \left(1 + \frac{1}{n}\right) \left(2 + \frac{1}{n}\right) \right] + 0 + 0 - 0 = 12 - \frac{9}{8}(1)(2) = 39/4$$

47.  $\Delta x = \frac{1}{n}$ ,  $x_k^* = \frac{2k-1}{2n}$

$$f(x_k^*)\Delta x = \frac{(2k-1)^2}{(2n)^2} \frac{1}{n} = \frac{k^2}{n^3} - \frac{k}{n^3} + \frac{1}{4n^3}$$

$$\sum_{k=1}^n f(x_k^*)\Delta x = \frac{1}{n^3} \sum_{k=1}^n k^2 - \frac{1}{n^3} \sum_{k=1}^n k + \frac{1}{4n^3} \sum_{k=1}^n 1$$

Using Theorem 5.4.4,

$$A = \lim_{n \rightarrow +\infty} \sum_{k=1}^n f(x_k^*)\Delta x = \frac{1}{3} + 0 + 0 = \frac{1}{3}$$

48.  $\Delta x = \frac{2}{n}$ ,  $x_k^* = -1 + \frac{2k-1}{n}$

$$f(x_k^*)\Delta x = \left( -1 + \frac{2k-1}{n} \right)^2 \frac{2}{n} = \frac{8k^2}{n^3} - \frac{8k}{n^3} + \frac{2}{n^3} - \frac{2}{n}$$

$$\sum_{k=1}^n f(x_k^* k) \Delta x = \frac{8}{n^3} \sum_{k=1}^n k^2 - \frac{8}{n^3} \sum_{k=1}^n k + \frac{2}{n^2} - 2$$

$$A = \lim_{n \rightarrow +\infty} \sum_{k=1}^n f(x_k^*) \Delta x = \frac{8}{3} + 0 + 0 - 2 = \frac{2}{3}$$

49.  $\Delta x = \frac{2}{n}, x_k^* = -1 + \frac{2k}{n}$

$$f(x_k^*) \Delta x = \left( -1 + \frac{2k}{n} \right) \frac{2}{n} = -\frac{2}{n} + 4 \frac{k}{n^2}$$

$$\sum_{k=1}^n f(x_k^*) \Delta x = -2 + \frac{4}{n^2} \sum_{k=1}^n k = -2 + \frac{4}{n^2} \frac{n(n+1)}{2} = -2 + 2 + \frac{2}{n}$$

$$A = \lim_{n \rightarrow +\infty} \sum_{k=1}^n f(x_k^*) \Delta x = 0$$

The area below the  $x$ -axis cancels the area above the  $x$ -axis.

50.  $\Delta x = \frac{3}{n}, x_k^* = -1 + \frac{3k}{n}$

$$f(x_k^*) \Delta x = \left( -1 + \frac{3k}{n} \right) \frac{3}{n} = -\frac{3}{n} + \frac{9}{n^2} k$$

$$\sum_{k=1}^n f(x_k^*) \Delta x = -3 + \frac{9}{n^2} \frac{n(n+1)}{2}$$

$$A = \lim_{n \rightarrow +\infty} \sum_{k=1}^n f(x_k^*) \Delta x = -3 + \frac{9}{2} + 0 = \frac{3}{2}$$

The area below the  $x$ -axis cancels the area above the  $x$ -axis that lies to the right of the line  $x = 1$ ; the remaining area is a trapezoid of width 1 and heights 1, 2, hence its area is  $\frac{1+2}{2} = \frac{3}{2}$

51.  $\Delta x = \frac{2}{n}, x_k^* = \frac{2k}{n}$

$$f(x_k^*) = \left[ \left( \frac{2k}{n} \right)^2 - 1 \right] \frac{2}{n} = \frac{8k^2}{n^3} - \frac{2}{n}$$

$$\sum_{k=1}^n f(x_k^*) \Delta x = \frac{8}{n^3} \sum_{k=1}^n k^2 - \frac{2}{n} \sum_{k=1}^n 1 = \frac{8}{n^3} \frac{n(n+1)(2n+1)}{6} - 2$$

$$A = \lim_{n \rightarrow +\infty} \sum_{k=1}^n f(x_k^*) \Delta x = \frac{16}{6} - 2 = \frac{2}{3}$$

52.  $\Delta x = \frac{2}{n}, x_k^* = -1 + \frac{2k}{n}$

$$f(x_k^*) \Delta x = \left( -1 + \frac{2k}{n} \right)^3 \frac{2}{n} = -\frac{2}{n} + 12 \frac{k}{n^2} - 24 \frac{k^2}{n^3} + 16 \frac{k^3}{n^4}$$

$$\sum_{k=1}^n f(x_k^*) \Delta x = -2 + \frac{12}{n^2} \frac{n(n+1)}{2} - \frac{24}{n^3} \frac{n(n+1)(2n+1)}{6} + \frac{16}{n^4} \left( \frac{n(n+1)}{2} \right)^2$$

$$A = \lim_{n \rightarrow +\infty} \sum_{k=1}^n f(x_k^*) = -2 + \frac{12}{2} - \frac{48}{6} + \frac{16}{2^2} = 0$$

53.  $\Delta x = \frac{b-a}{n}$ ,  $x_k^* = a + \frac{b-a}{n}(k-1)$

$$f(x_k^*)\Delta x = mx_k^*\Delta x = m \left[ a + \frac{b-a}{n}(k-1) \right] \frac{b-a}{n} = m(b-a) \left[ \frac{a}{n} + \frac{b-a}{n^2}(k-1) \right]$$

$$\sum_{k=1}^n f(x_k^*)\Delta x = m(b-a) \left[ a + \frac{b-a}{2} \cdot \frac{n-1}{n} \right]$$

$$A = \lim_{n \rightarrow +\infty} m(b-a) \left[ a + \frac{b-a}{2} \left( 1 - \frac{1}{n} \right) \right] = m(b-a) \frac{b+a}{2} = \frac{1}{2}m(b^2 - a^2)$$

54.  $\Delta x = \frac{b-a}{n}$ ,  $x_k^* = a + \frac{k}{n}(b-a)$

$$f(x_k^*)\Delta x = \frac{ma}{n}(b-a) + \frac{mk}{n^2}(b-a)^2$$

$$\sum_{k=1}^n f(x_k^*)\Delta x = ma(b-a) + \frac{m}{n^2}(b-a)^2 \frac{n(n+1)}{2}$$

$$A = \lim_{n \rightarrow +\infty} \sum_{k=1}^n f(x_k^*)\Delta x = ma(b-a) + \frac{m}{2}(b-a)^2 = m(b-a) \frac{a+b}{2}$$

55. (a) With  $x_k^*$  as the right endpoint,  $\Delta x = \frac{b}{n}$ ,  $x_k^* = \frac{b}{n}k$

$$f(x_k^*)\Delta x = (x_k^*)^3\Delta x = \frac{b^4}{n^4}k^3, \sum_{k=1}^n f(x_k^*)\Delta x = \frac{b^4}{n^4} \sum_{k=1}^n k^3 = \frac{b^4}{4} \frac{(n+1)^2}{n^2}$$

$$A = \lim_{n \rightarrow +\infty} \frac{b^4}{4} \left( 1 + \frac{1}{n} \right)^2 = b^4/4$$

(b)  $\Delta x = \frac{b-a}{n}$ ,  $x_k^* = a + \frac{b-a}{n}k$

$$\begin{aligned} f(x_k^*)\Delta x &= (x_k^*)^3\Delta x = \left[ a + \frac{b-a}{n}k \right]^3 \frac{b-a}{n} \\ &= \frac{b-a}{n} \left[ a^3 + \frac{3a^2(b-a)}{n}k + \frac{3a(b-a)^2}{n^2}k^2 + \frac{(b-a)^3}{n^3}k^3 \right] \end{aligned}$$

$$\begin{aligned} \sum_{k=1}^n f(x_k^*)\Delta x &= (b-a) \left[ a^3 + \frac{3}{2}a^2(b-a)\frac{n+1}{n} + \frac{1}{2}a(b-a)^2\frac{(n+1)(2n+1)}{n^2} \right. \\ &\quad \left. + \frac{1}{4}(b-a)^3\frac{(n+1)^2}{n^2} \right] \end{aligned}$$

$$\begin{aligned} A &= \lim_{n \rightarrow +\infty} \sum_{k=1}^n f(x_k^*)\Delta x \\ &= (b-a) \left[ a^3 + \frac{3}{2}a^2(b-a) + a(b-a)^2 + \frac{1}{4}(b-a)^3 \right] = \frac{1}{4}(b^4 - a^4). \end{aligned}$$

56. Let  $A$  be the area of the region under the curve and above the interval  $0 \leq x \leq 1$  on the  $x$ -axis, and let  $B$  be the area of the region between the curve and the interval  $0 \leq y \leq 1$  on the  $y$ -axis. Together  $A$  and  $B$  form the square of side 1, so  $A + B = 1$ .

But  $B$  can also be considered as the area between the curve  $x = y^2$  and the interval  $0 \leq y \leq 1$  on the  $y$ -axis. By Exercise 47 above,  $B = \frac{1}{3}$ , so  $A = 1 - \frac{1}{3} = \frac{2}{3}$ .

57. If  $n = 2m$  then  $2m + 2(m-1) + \cdots + 2 \cdot 2 + 2 = 2 \sum_{k=1}^m k = 2 \cdot \frac{m(m+1)}{2} = m(m+1) = \frac{n^2 + 2n}{4}$ ;

if  $n = 2m+1$  then  $(2m+1) + (2m-1) + \cdots + 5 + 3 + 1 = \sum_{k=1}^{m+1} (2k-1)$

$$= 2 \sum_{k=1}^{m+1} k - \sum_{k=1}^{m+1} 1 = 2 \cdot \frac{(m+1)(m+2)}{2} - (m+1) = (m+1)^2 = \frac{n^2 + 2n + 1}{4}$$

58.  $50 \cdot 30 + 49 \cdot 29 + \cdots + 22 \cdot 2 + 21 \cdot 1 = \sum_{k=1}^{30} k(k+20) = \sum_{k=1}^{30} k^2 + 20 \sum_{k=1}^{30} k = \frac{30 \cdot 31 \cdot 61}{6} + 20 \frac{30 \cdot 31}{2} = 18,755$

59. both are valid

60. none is valid

61.  $\sum_{k=1}^n (a_k - b_k) = (a_1 - b_1) + (a_2 - b_2) + \cdots + (a_n - b_n)$   
 $= (a_1 + a_2 + \cdots + a_n) - (b_1 + b_2 + \cdots + b_n) = \sum_{k=1}^n a_k - \sum_{k=1}^n b_k$

62.  $\sum_{k=1}^n [(k+1)^4 - k^4] = (n+1)^4 - 1$  (telescoping sum), expand the

quantity in brackets to get  $\sum_{k=1}^n (4k^3 + 6k^2 + 4k + 1) = (n+1)^4 - 1$ ,

$$4 \sum_{k=1}^n k^3 + 6 \sum_{k=1}^n k^2 + 4 \sum_{k=1}^n k + \sum_{k=1}^n 1 = (n+1)^4 - 1$$

$$\sum_{k=1}^n k^3 = \frac{1}{4} \left[ (n+1)^4 - 1 - 6 \sum_{k=1}^n k^2 - 4 \sum_{k=1}^n k - \sum_{k=1}^n 1 \right]$$

$$= \frac{1}{4} [(n+1)^4 - 1 - n(n+1)(2n+1) - 2n(n+1) - n]$$

$$= \frac{1}{4} (n+1)[(n+1)^3 - n(2n+1) - 2n - 1]$$

$$= \frac{1}{4} (n+1)(n^3 + n^2) = \frac{1}{4} n^2 (n+1)^2$$

63. (a)  $\sum_{k=1}^n 1$  means add 1 to itself  $n$  times, which gives the result.

(b)  $\frac{1}{n^2} \sum_{k=1}^n k = \frac{1}{n^2} \frac{n(n+1)}{2} = \frac{1}{2} + \frac{1}{2n}$ , so  $\lim_{n \rightarrow +\infty} \frac{1}{n^2} \sum_{k=1}^n k = \frac{1}{2}$

(c)  $\frac{1}{n^3} \sum_{k=1}^n k^2 = \frac{1}{n^3} \frac{n(n+1)(2n+1)}{6} = \frac{2}{6} + \frac{3}{6n} + \frac{1}{6n^2}$ , so  $\lim_{n \rightarrow +\infty} \frac{1}{n^3} \sum_{k=1}^n k^2 = \frac{1}{3}$

(d)  $\frac{1}{n^4} \sum_{k=1}^n k^3 = \frac{1}{n^4} \left( \frac{n(n+1)}{2} \right)^2 = \frac{1}{4} + \frac{1}{2n} + \frac{1}{4n^2}$ , so  $\lim_{n \rightarrow +\infty} \frac{1}{n^4} \sum_{k=1}^n k^3 = \frac{1}{4}$

## EXERCISE SET 5.5

1. (a)  $(4/3)(1) + (5/2)(1) + (4)(2) = 71/6$  (b) 2

2. (a)  $(\sqrt{2}/2)(\pi/2) + (-1)(3\pi/4) + (0)(\pi/2) + (\sqrt{2}/2)(\pi/4) = 3(\sqrt{2} - 2)\pi/8$   
 (b)  $3\pi/4$

3. (a)  $(-9/4)(1) + (3)(2) + (63/16)(1) + (-5)(3) = -117/16$   
 (b) 3

4. (a)  $(-8)(2) + (0)(1) + (0)(1) + (8)(2) = 0$  (b) 2

5.  $\int_{-1}^2 x^2 dx$

6.  $\int_1^2 x^3 dx$

7.  $\int_{-3}^3 4x(1 - 3x)dx$

8.  $\int_0^{\pi/2} \sin^2 x dx$



9. (a)  $\lim_{\max \Delta x_k \rightarrow 0} \sum_{k=1}^n 2x_k^* \Delta x_k; a = 1, b = 2$  (b)  $\lim_{\max \Delta x_k \rightarrow 0} \sum_{k=1}^n \frac{x_k^*}{x_k^* + 1} \Delta x_k; a = 0, b = 1$

10. (a)  $\lim_{\max \Delta x_k \rightarrow 0} \sum_{k=1}^n \sqrt{x_k^*} \Delta x_k, a = 1, b = 2$



(b)  $\lim_{\max \Delta x_k \rightarrow 0} \sum_{k=1}^n (1 + \cos x_k^*) \Delta x_k, a = -\pi/2, b = \pi/2$

11. (a)  $A = \frac{1}{2}(3)(3) = 9/2$


(b)  $-A = -\frac{1}{2}(1)(1+2) = -3/2$




(c)  $-A_1 + A_2 = -\frac{1}{2} + 8 = 15/2$




12. (a)  $A = \frac{1}{2}(1)(2) = 1$




(c)  $-A = -\frac{1}{2}(1/2)(1) = -1/4$




(b)  $A = \frac{1}{2}(2)(3/2 + 1/2) = 2$




(d)  $A_1 - A_2 = 1 - 1/4 = 3/4$




13. (a)  $A = 2(5) = 10$




(c)  $A_1 + A_2 = \frac{1}{2}(5)(5/2) + \frac{1}{2}(1)(1/2)$   
 $= 13/2$




(b) 0;  $A_1 = A_2$  by symmetry




(d)  $\frac{1}{2}[\pi(1)^2] = \pi/2$




14. (a)  $A = (6)(5) = 30$




(c)  $A_1 + A_2 = \frac{1}{2}(2)(2) + \frac{1}{2}(1)(1) = 5/2$



(b)  $-A_1 + A_2 = 0$  because  
 $A_1 = A_2$  by symmetry



(d)  $\frac{1}{4}\pi(2)^2 = \pi$



15. (a) 0.8

(b) -2.6

(c) -1.8

(d) -0.3

16. (a)  $\int_0^1 f(x)dx = \int_0^1 2xdx = x^2 \Big|_0^1 = 1$

(b)  $\int_{-1}^1 f(x)dx = \int_{-1}^1 2xdx = x^2 \Big|_{-1}^1 = 1^2 - (-1)^2 = 0$

(c)  $\int_1^{10} f(x)dx = \int_1^{10} 2dx = 2x \Big|_1^{10} = 18$

(d)  $\int_{1/2}^5 f(x)dx = \int_{1/2}^1 2xdx + \int_1^5 2dx = x^2 \Big|_{1/2}^1 + 2x \Big|_1^5 = 1^2 - (1/2)^2 + 2 \cdot 5 - 2 \cdot 1 = 3/4 + 8 = 35/4$

17.  $\int_{-1}^2 f(x)dx + 2 \int_{-1}^2 g(x)dx = 5 + 2(-3) = -1$

18.  $3 \int_1^4 f(x)dx - \int_1^4 g(x)dx = 3(2) - 10 = -4$

19.  $\int_1^5 f(x)dx = \int_0^5 f(x)dx - \int_0^1 f(x)dx = 1 - (-2) = 3$

20.  $\int_3^{-2} f(x)dx = - \int_{-2}^3 f(x)dx = - \left[ \int_{-2}^1 f(x)dx + \int_1^3 f(x)dx \right] = -(2 - 6) = 4$

21. (a)  $\int_0^1 xdx + 2 \int_0^1 \sqrt{1-x^2}dx = 1/2 + 2(\pi/4) = (1+\pi)/2$

(b)  $4 \int_{-1}^3 dx - 5 \int_{-1}^3 xdx = 4 \cdot 4 - 5(-1/2 + (3 \cdot 3)/2) = -4$

22. (a)  $\int_{-3}^0 2dx + \int_{-3}^0 \sqrt{9-x^2}dx = 2 \cdot 3 + (\pi(3)^2)/4 = 6 + 9\pi/4$

(b)  $\int_{-2}^2 dx - 3 \int_{-2}^2 |x|dx = 4 \cdot 1 - 3(2)(2 \cdot 2)/2 = -8$

23. (a)  $\sqrt{x} > 0$ ,  $1-x < 0$  on  $[2, 3]$  so the integral is negative(b)  $x^2 > 0$ ,  $3-\cos x > 0$  for all  $x$  so the integral is positive24. (a)  $x^4 > 0$ ,  $\sqrt{3-x} > 0$  on  $[-3, -1]$  so the integral is positive(b)  $x^3 - 9 < 0$ ,  $|x| + 1 > 0$  on  $[-2, 2]$  so the integral is negative

25.  $\int_0^{10} \sqrt{25-(x-5)^2}dx = \pi(5)^2/2 = 25\pi/2$

26.  $\int_0^3 \sqrt{9-(x-3)^2}dx = \pi(3)^2/4 = 9\pi/4$

27.  $\int_0^1 (3x+1)dx = 5/2$

28.  $\int_{-2}^2 \sqrt{4-x^2}dx = \pi(2)^2/2 = 2\pi$

29. (a)  $f$  is continuous on  $[-1, 1]$  so  $f$  is integrable there by Part (a) of Theorem 5.5.8
- (b)  $|f(x)| \leq 1$  so  $f$  is bounded on  $[-1, 1]$ , and  $f$  has one point of discontinuity, so by Part (b) of Theorem 5.5.8  $f$  is integrable on  $[-1, 1]$
- (c)  $f$  is not bounded on  $[-1, 1]$  because  $\lim_{x \rightarrow 0} f(x) = +\infty$ , so  $f$  is not integrable on  $[0, 1]$
- (d)  $f(x)$  is discontinuous at the point  $x = 0$  because  $\lim_{x \rightarrow 0} \frac{1}{x}$  does not exist.  $f$  is continuous elsewhere.  $-1 \leq f(x) \leq 1$  for  $x$  in  $[-1, 1]$  so  $f$  is bounded there. By Part (b), Theorem 5.5.8,  $f$  is integrable on  $[-1, 1]$ .
30. Each subinterval of a partition of  $[a, b]$  contains both rational and irrational numbers. If all  $x_k^*$  are chosen to be rational then
- $$\sum_{k=1}^n f(x_k^*) \Delta x_k = \sum_{k=1}^n (1) \Delta x_k = \sum_{k=1}^n \Delta x_k = b - a \text{ so } \lim_{\max \Delta x_k \rightarrow 0} \sum_{k=1}^n f(x_k^*) \Delta x_k = b - a.$$
- If all  $x_k^*$  are irrational then  $\lim_{\max \Delta x_k \rightarrow 0} \sum_{k=1}^n f(x_k^*) \Delta x_k = 0$ . Thus  $f$  is not integrable on  $[a, b]$  because the preceding limits are not equal.
31. (a) Let  $S_n = \sum_{k=1}^n f(x_k^*) \Delta x_k$  and  $S = \int_a^b f(x) dx$  then  $\sum_{k=1}^n c f(x_k^*) \Delta x_k = c S_n$  and we want to prove that  $\lim_{\max \Delta x_k \rightarrow 0} c S_n = c S$ . If  $c = 0$  the result follows immediately, so suppose that  $c \neq 0$  then for any  $\epsilon > 0$ ,  $|c S_n - c S| = |c||S_n - S| < \epsilon$  if  $|S_n - S| < \epsilon/|c|$ . But because  $f$  is integrable on  $[a, b]$ , there is a number  $\delta > 0$  such that  $|S_n - S| < \epsilon/|c|$  whenever  $\max \Delta x_k < \delta$  so  $|c S_n - c S| < \epsilon$  and hence  $\lim_{\max \Delta x_k \rightarrow 0} c S_n = c S$ .
- (b) Let  $R_n = \sum_{k=1}^n f(x_k^*) \Delta x_k$ ,  $S_n = \sum_{k=1}^n g(x_k^*) \Delta x_k$ ,  $T_n = \sum_{k=1}^n [f(x_k^*) + g(x_k^*)] \Delta x_k$ ,  $R = \int_a^b f(x) dx$ , and  $S = \int_a^b g(x) dx$  then  $T_n = R_n + S_n$  and we want to prove that  $\lim_{\max \Delta x_k \rightarrow 0} T_n = R + S$ .  $|T_n - (R + S)| = |(R_n - R) + (S_n - S)| \leq |R_n - R| + |S_n - S|$  so for any  $\epsilon > 0$   $|T_n - (R + S)| < \epsilon$  if  $|R_n - R| + |S_n - S| < \epsilon$ . Because  $f$  and  $g$  are integrable on  $[a, b]$ , there are numbers  $\delta_1$  and  $\delta_2$  such that  $|R_n - R| < \epsilon/2$  for  $\max \Delta x_k < \delta_1$  and  $|S_n - S| < \epsilon/2$  for  $\max \Delta x_k < \delta_2$ . If  $\delta = \min(\delta_1, \delta_2)$  then  $|R_n - R| < \epsilon/2$  and  $|S_n - S| < \epsilon/2$  for  $\max \Delta x_k < \delta$  thus  $|R_n - R| + |S_n - S| < \epsilon$  and so  $|T_n - (R + S)| < \epsilon$  for  $\max \Delta x_k < \delta$  which shows that  $\lim_{\max \Delta x_k \rightarrow 0} T_n = R + S$ .
32. For the smallest, find  $x_k^*$  so that  $f(x_k^*)$  is minimum on each subinterval:  $x_1^* = 1$ ,  $x_2^* = 3/2$ ,  $x_3^* = 3$  so  $(2)(1) + (7/4)(2) + (4)(1) = 9.5$ . For the largest, find  $x_k^*$  so that  $f(x_k^*)$  is maximum on each subinterval:  $x_1^* = 0$ ,  $x_2^* = 3$ ,  $x_3^* = 4$  so  $(4)(1) + (4)(2) + (8)(1) = 20$ .

33.  $\Delta x_k = \frac{4k^2}{n^2} - \frac{4(k-1)^2}{n^2} = \frac{4}{n^2}(2k-1)$ ,  $x_k^* = \frac{4k^2}{n^2}$ ,

$$f(x_k^*) = \frac{2k}{n}, f(x_k^*) \Delta x_k = \frac{8k}{n^3}(2k-1) = \frac{8}{n^3}(2k^2 - k),$$

$$\sum_{k=1}^n f(x_k^*) \Delta x_k = \frac{8}{n^3} \sum_{k=1}^n (2k^2 - k) = \frac{8}{n^3} \left[ \frac{1}{3} n(n+1)(2n+1) - \frac{1}{2} n(n+1) \right] = \frac{4}{3} \frac{(n+1)(4n-1)}{n^2},$$

$$\lim_{n \rightarrow +\infty} \sum_{k=1}^n f(x_k^*) \Delta x_k = \lim_{n \rightarrow +\infty} \frac{4}{3} \left( 1 + \frac{1}{n} \right) \left( 4 - \frac{1}{n} \right) = \frac{16}{3}.$$

- 34.** For any partition of  $[a, b]$  use the right endpoints to form the sum  $\sum_{k=1}^n f(x_k^*) \Delta x_k$ . Since  $f(x_k^*) = 0$  for each  $k$ , the sum is zero and so is  $\int_a^b f(x) dx = \lim_{n \rightarrow +\infty} \sum_{k=1}^n f(x_k^*) \Delta x_k$ .
- 35.** With  $f(x) = g(x)$  then  $f(x) - g(x) = 0$  for  $a < x \leq b$ . By Theorem 5.5.4(b)  
 $\int_a^b f(x) dx = \int_a^b [(f(x) - g(x) + g(x)] dx = \int_a^b [f(x) - g(x)] dx + \int_a^b g(x) dx$ .  
But the first term on the right hand side is zero (from Exercise 34), so  
 $\int_a^b f(x) dx = \int_a^b g(x) dx$
- 36.** Choose any large positive integer  $N$  and any partition of  $[0, a]$ . Then choose  $x_1^*$  in the first interval so small that  $f(x_1^*) \Delta x_1 > N$ . For example choose  $x_1^* < \Delta x_1/N$ . Then with this partition and choice of  $x_1^*$ ,  $\sum_{k=1}^n f(x_k^*) \Delta x_k > f(x_1^*) \Delta x_1 > N$ . This shows that the sum is dependent on partition and/or points, so Definition 5.5.1 is not satisfied.

## EXERCISE SET 5.6

1. (a)  $\int_0^2 (2-x) dx = (2x - x^2/2) \Big|_0^2 = 4 - 4/2 = 2$
1. (b)  $\int_{-1}^1 2dx = 2x \Big|_{-1}^1 = 2(1) - 2(-1) = 4$
1. (c)  $\int_1^3 (x+1) dx = (x^2/2 + x) \Big|_1^3 = 9/2 + 3 - (1/2 + 1) = 6$
2. (a)  $\int_0^5 x dx = x^2/2 \Big|_0^5 = 25/2$
2. (b)  $\int_3^9 5dx = 5x \Big|_3^9 = 5(9) - 5(3) = 30$
2. (c)  $\int_{-1}^2 (x+3) dx = (x^2/2 + 3x) \Big|_{-1}^2 = 4/2 + 6 - (1/2 - 3) = 21/2$
3.  $\int_2^3 x^3 dx = x^4/4 \Big|_2^3 = 81/4 - 16/4 = 65/4$
4.  $\int_{-1}^1 x^4 dx = x^5/5 \Big|_{-1}^1 = 1/5 - (-1)/5 = 2/5$
5.  $\int_1^9 \sqrt{x} dx = \frac{2}{3} x^{3/2} \Big|_1^9 = \frac{2}{3} (27 - 1) = 52/3$
6.  $\int_1^4 x^{-3/5} dx = \frac{5}{2} x^{2/5} \Big|_1^4 = \frac{5}{2} (4^{2/5} - 1)$
7.  $\left( \frac{1}{3} x^3 - 2x^2 + 7x \right) \Big|_{-3}^0 = 48$
8.  $\left( \frac{1}{2} x^2 + \frac{1}{5} x^5 \right) \Big|_{-1}^2 = 81/10$
9.  $\int_1^3 x^{-2} dx = -\frac{1}{x} \Big|_1^3 = 2/3$
10.  $\int_1^2 x^{-6} dx = -\frac{1}{5x^5} \Big|_1^2 = 31/160$

11.  $\left[ \frac{4}{5}x^{5/2} \right]_4^9 = 844/5$

12.  $\left( 3x^{5/3} + \frac{4}{x} \right) \Big|_1^8 = 179/2$

13.  $-\cos \theta \Big|_{-\pi/2}^{\pi/2} = 0$

14.  $\tan \theta \Big|_0^{\pi/4} = 1$

15.  $\sin x \Big|_{-\pi/4}^{\pi/4} = \sqrt{2}$

16.  $\left( \frac{1}{2}x^2 - \sec x \right) \Big|_0^1 = 3/2 - \sec(1)$

17.  $\left( 6\sqrt{t} - \frac{10}{3}t^{3/2} + \frac{2}{\sqrt{t}} \right) \Big|_1^4 = -55/3$

18.  $\left( 8\sqrt{y} + \frac{4}{3}y^{3/2} - \frac{2}{3y^{3/2}} \right) \Big|_4^9 = 10819/324$

19.  $\left( \frac{1}{2}x^2 - 2 \cot x \right) \Big|_{\pi/6}^{\pi/2} = \pi^2/9 + 2\sqrt{3}$

20.  $\left( a^{1/2}x - \frac{2}{3}x^{3/2} \right) \Big|_a^{4a} = -\frac{5}{3}a^{3/2}$

21. (a)  $\int_0^{3/2} (3 - 2x)dx + \int_{3/2}^2 (2x - 3)dx = (3x - x^2) \Big|_0^{3/2} + (x^2 - 3x) \Big|_{3/2}^2 = 9/4 + 1/4 = 5/2$

(b)  $\int_0^{\pi/2} \cos x dx + \int_{\pi/2}^{3\pi/4} (-\cos x)dx = \sin x \Big|_0^{\pi/2} - \sin x \Big|_{\pi/2}^{3\pi/4} = 2 - \sqrt{2}/2$

22. (a)  $\int_{-1}^0 \sqrt{2-x} dx + \int_0^2 \sqrt{2+x} dx = -\frac{2}{3}(2-x)^{3/2} \Big|_{-1}^0 + \frac{2}{3}(2+x)^{3/2} \Big|_0^2 = -\frac{2}{3}(2\sqrt{2} - 3\sqrt{3}) + \frac{2}{3}(8 - 2\sqrt{2}) = \frac{2}{3}(8 - 4\sqrt{2} + 3\sqrt{3})$

(b)  $\int_0^{\pi/6} (1/2 - \sin x) dx + \int_{\pi/6}^{\pi/2} (\sin x - 1/2) dx = (x/2 + \cos x) \Big|_0^{\pi/6} - (\cos x + x/2) \Big|_{\pi/6}^{\pi/2} = (\pi/12 + \sqrt{3}/2) - 1 - \pi/4 + (\sqrt{3}/2 + \pi/12) = \sqrt{3} - \pi/12 - 1$

23. (a) 17/6

(b)  $F(x) = \begin{cases} \frac{1}{2}x^2, & x \leq 1 \\ \frac{1}{3}x^3 + \frac{1}{6}, & x > 1 \end{cases}$

24. (a)  $\int_0^1 \sqrt{x} dx + \int_1^4 \frac{1}{x^2} dx = \frac{2}{3}x^{3/2} \Big|_0^1 - \frac{1}{x} \Big|_1^4 = 17/12$

(b)  $F(x) = \begin{cases} \frac{2}{3}x^{3/2}, & x < 1 \\ -\frac{1}{x} + \frac{5}{3}, & x \geq 1 \end{cases}$

25. 0.665867079;  $\int_1^3 \frac{1}{x^2} dx = -\frac{1}{x} \Big|_1^3 = 2/3$

26. 1.000257067;  $\int_0^{\pi/2} \sin x dx = -\cos x \Big|_0^{\pi/2} = 1$

27.  $3.106017890; \int_{-1}^1 \sec^2 x dx = \tan x \Big|_{-1}^1 = 2 \tan 1 \approx 3.114815450$

29.  $A = \int_0^3 (x^2 + 1)dx = \left( \frac{1}{3}x^3 + x \right) \Big|_0^3 = 12$


30.  $A = \int_1^2 (-x^2 + 3x - 2)dx = \left( -\frac{1}{3}x^3 + \frac{3}{2}x^2 - 2x \right) \Big|_1^2 = 1/6$

31.  $A = \int_0^{2\pi/3} 3 \sin x dx = -3 \cos x \Big|_0^{2\pi/3} = 9/2$       32.  $A = - \int_{-2}^{-1} x^3 dx = -\frac{1}{4}x^4 \Big|_{-2}^{-1} = 15/4$

33.  $A_1 = \int_{-3}^{-2} (x^2 - 3x - 10)dx = \left( \frac{1}{3}x^3 - \frac{3}{2}x^2 - 10x \right) \Big|_{-3}^{-2} = 23/6,$

$$A_2 = - \int_{-2}^5 (x^2 - 3x - 10)dx = 343/6,$$

$$A_3 = \int_5^8 (x^2 - 3x - 10)dx = 243/6, A = A_1 + A_2 + A_3 = 203/2$$



34. (a) the area is positive

(b)  $\int_{-2}^5 \left( \frac{1}{100}x^3 - \frac{1}{20}x^2 - \frac{1}{25}x + \frac{1}{5} \right) dx = \left( \frac{1}{400}x^4 - \frac{1}{60}x^3 - \frac{1}{50}x^2 + \frac{1}{5}x \right) \Big|_{-2}^5 = \frac{343}{1200}$

35. (a) the area between the curve and the  $x$ -axis breaks into equal parts, one above and one below the  $x$ -axis, so the integral is zero

(b)  $\int_{-1}^1 x^3 dx = \frac{1}{4}x^4 \Big|_{-1}^1 = \frac{1}{4}(1^4 - (-1)^4) = 0;$

$$\int_{-\pi/2}^{\pi/2} \sin x dx = -\cos x \Big|_{-\pi/2}^{\pi/2} = -\cos(\pi/2) + \cos(-\pi/2) = 0 + 0 = 0$$

(c) The area on the left side of the  $y$ -axis is equal to the area on the right side, so

$$\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$$

(d)  $\int_{-1}^1 x^2 dx = \frac{1}{3}x^3 \Big|_{-1}^1 = \frac{1}{3}(1^3 - (-1)^3) = \frac{2}{3} = 2 \int_0^1 x^2 dx;$

$$\int_{-\pi/2}^{\pi/2} \cos x dx = \sin x \Big|_{-\pi/2}^{\pi/2} = \sin(\pi/2) - \sin(-\pi/2) = 1 + 1 = 2 = 2 \int_0^{\pi/2} \cos x dx$$

36. The numerator is an odd function and the denominator is an even function, so the integrand is an odd function and the integral is zero.

37. (a)  $x^3 + 1$

(b)  $F(x) = \left( \frac{1}{4}t^4 + t \right) \Big|_1^x = \frac{1}{4}x^4 + x - \frac{5}{4}; F'(x) = x^3 + 1$

38. (a)  $\cos 2x$

(b)  $F(x) = \frac{1}{2} \sin 2t \Big|_{\pi/4}^x = \frac{1}{2} \sin 2x - \frac{1}{2}, F'(x) = \cos 2x$

39. (a)  $\sin \sqrt{x}$       (b)  $\sqrt{1 + \cos^2 x}$

40. (a)  $\frac{1}{1 + \sqrt{x}}$       (b)  $\frac{1}{1 + x + x^2}$

41.  $-\frac{x}{\cos x}$

42.  $|u|$

43.  $F'(x) = \sqrt{3x^2 + 1}$ ,  $F''(x) = \frac{3x}{\sqrt{3x^2 + 1}}$

(a) 0

(b)  $\sqrt{13}$

(c)  $6/\sqrt{13}$

44.  $F'(x) = \frac{\cos x}{x^2 + 3}$ ,  $F''(x) = \frac{-(x^2 + 3)\sin x - 2x\cos x}{(x^2 + 3)^2}$

(a) 0

(b)  $1/3$


(c) 0

45. (a)  $F'(x) = \frac{x-3}{x^2+7} = 0$  when  $x = 3$ , which is a relative minimum, and hence the absolute minimum, by the first derivative test.

(b) increasing on  $[3, +\infty)$ , decreasing on  $(-\infty, 3]$

(c)  $F''(x) = \frac{7+6x-x^2}{(x^2+7)^2} = \frac{(7-x)(1+x)}{(x^2+7)^2}$ ; concave up on  $(-1, 7)$ , concave down on  $(-\infty, -1)$  and on  $(7, +\infty)$

46.



47. (a)  $(0, +\infty)$  because  $f$  is continuous there and 1 is in  $(0, +\infty)$

(b) at  $x = 1$  because  $F(1) = 0$

48. (a)  $(-3, 3)$  because  $f$  is continuous there and 1 is in  $(-3, 3)$

(b) at  $x = 1$  because  $F(1) = 0$

49. (a)  $f_{\text{ave}} = \frac{1}{9} \int_0^9 x^{1/2} dx = 2$ ;  $\sqrt{x^*} = 2$ ,  $x^* = 4$

(b)  $f_{\text{ave}} = \frac{1}{3} \int_{-1}^2 (3x^2 + 2x + 1) dx = \left. \frac{1}{3}(x^3 + x^2 + x) \right|_{-1}^2 = 5$ ;  $3x^{*2} + 2x^* + 1 = 5$ ,

with solutions  $x^* = -(1/3)(1 \pm \sqrt{13})$ , but only  $x^* = -(1/3)(1 - \sqrt{13})$  lies in the interval  $[-1, 2]$ .

50. (a)  $f_{\text{ave}} = \frac{1}{2\pi} \int_{-\pi}^{\pi} \sin x dx = 0$ ;  $\sin x^* = 0$ ,  $x^* = -\pi, 0, \pi$

(b)  $f_{\text{ave}} = \frac{1}{2} \int_1^3 \frac{1}{x^2} dx = \frac{1}{3}$ ;  $\frac{1}{(x^*)^2} = \frac{1}{3}$ ,  $x^* = \sqrt{3}$

51.  $\sqrt{2} \leq \sqrt{x^3 + 2} \leq \sqrt{29}$ , so  $3\sqrt{2} \leq \int_0^3 \sqrt{x^3 + 2} dx \leq 3\sqrt{29}$

52. Let  $f(x) = x \sin x$ ,  $f(0) = f(1) = 0$ ,  $f'(x) = \sin x + x \cos x = 0$  when  $x = -\tan x$ ,  $x \approx 2.0288$ , so  $f$  has an absolute maximum at  $x \approx 2.0288$ ;  $f(2.0288) \approx 1.8197$ , so  $0 \leq x \sin x \leq 1.82$  and  $0 \leq \int_0^\pi x \sin x dx \leq 1.82\pi = 5.72$


53. (a)  $[cF(x)]_a^b = cF(b) - cF(a) = c[F(b) - F(a)] = c [F(x)]_a^b$   
 (b)  $[F(x) + G(x)]_a^b = [F(b) + G(b)] - [F(a) + G(a)]$   
 $= [F(b) - F(a)] + [G(b) - G(a)] = F(x)]_a^b + G(x)]_a^b$   
 (c)  $[F(x) - G(x)]_a^b = [F(b) - G(b)] - [F(a) - G(a)]$   
 $= [F(b) - F(a)] - [G(b) - G(a)] = F(x)]_a^b - G(x)]_a^b$

54. Let  $f$  be continuous on a closed interval  $[a, b]$  and let  $F$  be an antiderivative of  $f$  on  $[a, b]$ . By Theorem 4.8.2,  $\frac{F(b) - F(a)}{b - a} = F'(x^*)$  for some  $x^*$  in  $(a, b)$ . By Theorem 5.6.1,  
 $\int_a^b f(x) dx = F(b) - F(a)$ , i.e.  $\int_a^b f(x) dx = F'(x^*)(b - a) = f(x^*)(b - a)$ .

## EXERCISE SET 5.7

1. (a) the increase in height in inches, during the first ten years  
 (b) the change in the radius in centimeters, during the time interval  $t = 1$  to  $t = 2$  seconds  
 (c) the change in the speed of sound in ft/s, during an increase in temperature from  $t = 32^\circ\text{F}$  to  $t = 100^\circ\text{F}$   
 (d) the displacement of the particle in cm, during the time interval  $t = t_1$  to  $t = t_2$  seconds
2. (a)  $\int_0^1 V(t) dt$  gal  
 (b) the change  $f(x_1) - f(x_2)$  in the values of  $f$  over the interval
3. (a) displ =  $s(3) - s(0)$   
 $= \int_0^3 v(t) dt = \int_0^2 (1-t) dt + \int_2^3 (t-3) dt = (t - t^2/2) \Big|_0^2 + (t^2/2 - 3t) \Big|_2^3 = -1/2;$   
 $\text{dist} = \int_0^3 |v(t)| dt = (t - t^2/2) \Big|_0^1 + (t^2/2 - t) \Big|_1^2 - (t^2/2 - 3t) \Big|_2^3 = 3/2$
- (b) displ =  $s(3) - s(0)$   
 $= \int_0^3 v(t) dt = \int_0^1 t dt + \int_1^2 dt + \int_2^3 (5-2t) dt = t^2/2 \Big|_0^1 + t \Big|_1^2 + (5t - t^2) \Big|_2^3 = 3/2;$   
 $\text{dist} = \int_0^1 t dt + \int_1^2 dt + \int_2^{5/2} (5-2t) dt + \int_{5/2}^3 (2t-5) dt$   
 $= t^2/2 \Big|_0^1 + t \Big|_1^2 + (5t - t^2) \Big|_2^{5/2} + (t^2 - 5t) \Big|_{5/2}^3 = 2$

4.



5. (a)  $v(t) = 20 + \int_0^t a(u)du$ ; add areas of the small blocks to get

$$v(4) \approx 20 + 1.4 + 3.0 + 4.7 + 6.2 = 35.3 \text{ m/s}$$

(b)  $v(6) = v(4) + \int_4^6 a(u)du \approx 35.3 + 7.5 + 8.6 = 51.4 \text{ m/s}$

6.  $a > 0$  and therefore (Theorem 5.5.6(a))  $v > 0$ , so the particle is always speeding up for  $0 < t < 10$

7. (a)  $s(t) = \int (t^3 - 2t^2 + 1)dt = \frac{1}{4}t^4 - \frac{2}{3}t^3 + t + C$ ,

$$s(0) = \frac{1}{4}(0)^4 - \frac{2}{3}(0)^3 + 0 + C = 1, C = 1, s(t) = \frac{1}{4}t^4 - \frac{2}{3}t^3 + t + 1$$

(b)  $v(t) = \int 4 \cos 2t dt = 2 \sin 2t + C_1, v(0) = 2 \sin 0 + C_1 = -1, C_1 = -1$ ,

$$v(t) = 2 \sin 2t - 1, s(t) = \int (2 \sin 2t - 1)dt = -\cos 2t - t + C_2,$$

$$s(0) = -\cos 0 - 0 + C_2 = -3, C_2 = -2, s(t) = -\cos 2t - t - 2$$

8. (a)  $s(t) = \int (1 + \sin t)dt = t - \cos t + C, s(0) = 0 - \cos 0 + C = -3, C = -2, s(t) = t - \cos t - 2$

(b)  $v(t) = \int (t^2 - 3t + 1)dt = \frac{1}{3}t^3 - \frac{3}{2}t^2 + t + C_1$ ,

$$v(0) = \frac{1}{3}(0)^3 - \frac{3}{2}(0)^2 + 0 + C_1 = 0, C_1 = 0, v(t) = \frac{1}{3}t^3 - \frac{3}{2}t^2 + t,$$

$$s(t) = \int \left( \frac{1}{3}t^3 - \frac{3}{2}t^2 + t \right) dt = \frac{1}{12}t^4 - \frac{1}{2}t^3 + \frac{1}{2}t^2 + C_2,$$

$$s(0) = \frac{1}{12}(0)^4 - \frac{1}{2}(0)^3 + \frac{1}{2}(0)^2 + C_2 = 0, C_2 = 0, s(t) = \frac{1}{12}t^4 - \frac{1}{2}t^3 + \frac{1}{2}t^2$$

9. (a)  $s(t) = \int (2t - 3)dt = t^2 - 3t + C, s(1) = (1)^2 - 3(1) + C = 5, C = 7, s(t) = t^2 - 3t + 7$

(b)  $v(t) = \int \cos t dt = \sin t + C_1, v(\pi/2) = 2 = 1 + C_1, C_1 = 1, v(t) = \sin t + 1$ ,

$$s(t) = \int (\sin t + 1)dt = -\cos t + t + C_2, s(\pi/2) = 0 = \pi/2 + C_2, C_2 = -\pi/2,$$

$$s(t) = -\cos t + t - \pi/2$$

10. (a)  $s(t) = \int t^{2/3} dt = \frac{3}{5}t^{5/3} + C, s(8) = 0 = \frac{3}{5}32 + C, C = -\frac{96}{5}, s(t) = \frac{3}{5}t^{5/3} - \frac{96}{5}$

(b)  $v(t) = \int \sqrt{t} dt = \frac{2}{3}t^{3/2} + C_1, v(4) = 1 = \frac{2}{3}8 + C_1, C_1 = -\frac{13}{3}, v(t) = \frac{2}{3}t^{3/2} - \frac{13}{3}$ ,

$$s(t) = \int \left( \frac{2}{3}t^{3/2} - \frac{13}{3} \right) dt = \frac{4}{15}t^{5/2} - \frac{13}{3}t + C_2, s(4) = -5 = \frac{4}{15}32 - \frac{13}{3}4 + C_2 = -\frac{44}{5} + C_2,$$

$$C_2 = \frac{19}{5}, s(t) = \frac{4}{15}t^{5/2} - \frac{13}{3}t + \frac{19}{5}$$

11. (a) displacement =  $s(\pi/2) - s(0) = \int_0^{\pi/2} \sin t dt = -\cos t \Big|_0^{\pi/2} = 1$  m

$$\text{distance} = \int_0^{\pi/2} |\sin t| dt = 1$$
 m

(b) displacement =  $s(2\pi) - s(\pi/2) = \int_{\pi/2}^{2\pi} \cos t dt = \sin t \Big|_{\pi/2}^{2\pi} = -1$  m

$$\text{distance} = \int_{\pi/2}^{2\pi} |\cos t| dt = - \int_{\pi/2}^{3\pi/2} \cos t dt + \int_{3\pi/2}^{2\pi} \cos t dt = 3$$
 m

12. (a) displacement =  $s(6) - s(0) = \int_0^6 (2t - 4) dt = (t^2 - 4t) \Big|_0^6 = 12$  m

$$\text{distance} = \int_0^6 |2t - 4| dt = \int_0^2 (4 - 2t) dt + \int_2^6 (2t - 4) dt = (4t - t^2) \Big|_0^2 + (t^2 - 4t) \Big|_2^6 = 20$$
 m

(b) displacement =  $\int_0^5 |t - 3| dt = \int_0^3 -(t - 3) dt + \int_3^5 (t - 3) dt = 13/2$  m

$$\text{distance} = \int_0^5 |t - 3| dt = 13/2$$
 m

13. (a)  $v(t) = t^3 - 3t^2 + 2t = t(t-1)(t-2)$

$$\text{displacement} = \int_0^3 (t^3 - 3t^2 + 2t) dt = 9/4$$
 m

$$\text{distance} = \int_0^3 |v(t)| dt = \int_0^1 v(t) dt + \int_1^2 -v(t) dt + \int_2^3 v(t) dt = 11/4$$
 m

(b) displacement =  $\int_0^3 (\sqrt{t} - 2) dt = 2\sqrt{3} - 6$  m

$$\text{distance} = \int_0^3 |v(t)| dt = - \int_0^3 v(t) dt = 6 - 2\sqrt{3}$$
 m

14. (a) displacement =  $\int_1^3 \left( \frac{1}{2} - \frac{1}{t^2} \right) dt = 1/3$  m

$$\text{distance} = \int_1^3 |v(t)| dt = - \int_1^{\sqrt{2}} v(t) dt + \int_{\sqrt{2}}^3 v(t) dt = 10/3 - 2\sqrt{2}$$
 m

(b) displacement =  $\int_4^9 3t^{-1/2} dt = 6$  m

$$\text{distance} = \int_4^9 |v(t)| dt = \int_4^9 v(t) dt = 6$$
 m

15.  $v(t) = -2t + 3$

$$\text{displacement} = \int_1^4 (-2t + 3) dt = -6$$
 m

$$\text{distance} = \int_1^4 |-2t + 3| dt = \int_1^{3/2} (-2t + 3) dt + \int_{3/2}^4 (2t - 3) dt = 13/2$$
 m

16.  $v(t) = \frac{1}{2}t^2 - 2t$

$$\text{displacement} = \int_1^5 \left( \frac{1}{2}t^2 - 2t \right) dt = -10/3 \text{ m}$$

$$\text{distance} = \int_1^5 \left| \frac{1}{2}t^2 - 2t \right| dt = \int_1^4 -\left( \frac{1}{2}t^2 - 2t \right) dt + \int_4^5 \left( \frac{1}{2}t^2 - 2t \right) dt = 17/3 \text{ m}$$

17.  $v(t) = \frac{2}{5}\sqrt{5t+1} + \frac{8}{5}$

$$\text{displacement} = \int_0^3 \left( \frac{2}{5}\sqrt{5t+1} + \frac{8}{5} \right) dt = \frac{4}{75}(5t+1)^{3/2} + \frac{8}{5}t \Big|_0^3 = 204/25 \text{ m}$$

$$\text{distance} = \int_0^3 |v(t)| dt = \int_0^3 v(t) dt = 204/25 \text{ m}$$

18.  $v(t) = -\cos t + 2$

$$\text{displacement} = \int_{\pi/4}^{\pi/2} (-\cos t + 2) dt = (\pi + \sqrt{2} - 2)/2 \text{ m}$$

$$\text{distance} = \int_{\pi/4}^{\pi/2} |- \cos t + 2| dt = \int_{\pi/4}^{\pi/2} (-\cos t + 2) dt = (\pi + \sqrt{2} - 2)/2 \text{ m}$$

19. (a)  $s = \int \sin \frac{1}{2}\pi t dt = -\frac{2}{\pi} \cos \frac{1}{2}\pi t + C$

$$s = 0 \text{ when } t = 0 \text{ which gives } C = \frac{2}{\pi} \text{ so } s = -\frac{2}{\pi} \cos \frac{1}{2}\pi t + \frac{2}{\pi}.$$

$$a = \frac{dv}{dt} = \frac{\pi}{2} \cos \frac{1}{2}\pi t. \text{ When } t = 1 : s = 2/\pi, v = 1, |v| = 1, a = 0.$$

(b)  $v = -3 \int t dt = -\frac{3}{2}t^2 + C_1, v = 0 \text{ when } t = 0 \text{ which gives } C_1 = 0 \text{ so } v = -\frac{3}{2}t^2$

$$s = -\frac{3}{2} \int t^2 dt = -\frac{1}{2}t^3 + C_2, s = 1 \text{ when } t = 0 \text{ which gives } C_2 = 1 \text{ so } s = -\frac{1}{2}t^3 + 1.$$

$$\text{When } t = 1 : s = 1/2, v = -3/2, |v| = 3/2, a = -3.$$

20. (a) negative, because  $v$  is decreasing

(b) speeding up when  $av > 0$ , so  $2 < t < 5$ ; slowing down when  $1 < t < 2$

(c) negative, because the area between the graph of  $v(t)$  and the t-axis appears to be greater where  $v < 0$  compared to where  $v > 0$




21.  $A = A_1 + A_2 = \int_0^1 (1 - x^2) dx + \int_1^3 (x^2 - 1) dx = 2/3 + 20/3 = 22/3$

22.  $A = A_1 + A_2 = \int_0^\pi \sin x dx - \int_\pi^{3\pi/2} \sin x dx = 2 + 1 = 3$




23.  $A = A_1 + A_2 = \int_{-1}^0 [1 - \sqrt{x+1}] dx + \int_0^1 [\sqrt{x+1} - 1] dx$   
 $= \left( x - \frac{2}{3}(x+1)^{3/2} \right) \Big|_{-1}^0 + \left( \frac{2}{3}(x+1)^{3/2} - x \right) \Big|_0^1 = -\frac{2}{3} + 1 + \frac{4\sqrt{2}}{3} - 1 - \frac{2}{3} = 4\frac{\sqrt{2}-1}{3}$

24.  $A = A_1 + A_2 = \int_{1/2}^1 \frac{1-x^2}{x^2} dx + \int_1^2 \frac{x^2-1}{x^2} dx = \left( -\frac{1}{x} - x \right) \Big|_{1/2}^1 + \left( x + \frac{1}{x} \right) \Big|_1^2$   
 $= -2 + 2 + \frac{1}{2} + 2 + \frac{1}{2} - 2 = 1$


25.  $s(t) = \frac{20}{3}t^3 - 50t^2 + 50t + s_0$ ,  $s(0) = 0$  gives  $s_0 = 0$ , so  $s(t) = \frac{20}{3}t^3 - 50t^2 + 50t$ ,  $a(t) = 40t - 100$



26.  $v(t) = 2t^2 - 30t + v_0$ ,  $v(0) = 3 = v_0$ , so  $v(t) = 2t^2 - 30t + 3$ ,  $s(t) = \frac{2}{3}t^3 - 15t^2 + 3t + s_0$ ,  
 $s(0) = -5 = s_0$ , so  $s(t) = \frac{2}{3}t^3 - 15t^2 + 3t - 5$




27. (a) From the graph the velocity is at first positive, but then turns negative, then positive again. The displacement, which is the cumulative area from  $x = 0$  to  $x = 5$ , starts positive, turns negative, and then turns positive again.  
(b) displ =  $5/2 - \sin 5 + 5 \cos 5$




28. (a) If  $t_0 < 1$  then the area between the velocity curve and the  $t$ -axis, between  $t = 0$  and  $t = t_0$ , will always be positive, so the displacement will be positive.


(b) displ =  $\frac{\pi^2 + 4}{2\pi^2}$



29. (a)  $a(t) = \begin{cases} 0, & t < 4 \\ -10, & t > 4 \end{cases}$



(b)  $v(t) = \begin{cases} 25, & t < 4 \\ 65 - 10t, & t > 4 \end{cases}$



(c)  $x(t) = \begin{cases} 25t, & t < 4 \\ 65t - 5t^2 - 80, & t > 4 \end{cases}$ , so  $x(8) = 120$ ,  $x(12) = -20$

(d)  $x(6.5) = 131.25$

30. (a) From (9)  $t = \frac{v - v_0}{a}$ ; from that and (8)

$$s - s_0 = v_0 \frac{v - v_0}{a} + \frac{1}{2} a \frac{(v - v_0)^2}{a^2}; \text{ multiply through by } a \text{ to get}$$

$$a(s - s_0) = v_0(v - v_0) + \frac{1}{2}(v - v_0)^2 = (v - v_0) \left[ v_0 + \frac{1}{2}(v - v_0) \right] = \frac{1}{2}(v^2 - v_0^2). \text{ Thus}$$

$$a = \frac{v^2 - v_0^2}{2(s - s_0)}.$$

(b) Put the last result of Part (a) into the first equation of Part (a) to obtain

$$t = \frac{v - v_0}{a} = (v - v_0) \frac{2(s - s_0)}{v^2 - v_0^2} = \frac{2(s - s_0)}{v + v_0}.$$

(c) From (9)  $v_0 = v - at$ ; use this in (8) to get

$$s - s_0 = (v - at)t + \frac{1}{2}at^2 = vt - \frac{1}{2}at^2$$

This expression contains no  $v_0$  terms and so differs from (8).

31. (a)  $a = -1 \text{ mi/h/s} = -22/15 \text{ ft/s}^2$

(b)  $a = 30 \text{ km/h/min} = 1/7200 \text{ km/s}^2$

32. Take  $t = 0$  when deceleration begins, then  $a = -10$  so  $v = -10t + C_1$ , but  $v = 88$  when  $t = 0$  which gives  $C_1 = 88$  thus  $v = -10t + 88$ ,  $t \geq 0$

(a)  $v = 45 \text{ mi/h} = 66 \text{ ft/s}$ ,  $66 = -10t + 88$ ,  $t = 2.2 \text{ s}$

(b)  $v = 0$  (the car is stopped) when  $t = 8.8 \text{ s}$

$$s = \int v dt = \int (-10t + 88) dt = -5t^2 + 88t + C_2, \text{ and taking } s = 0 \text{ when } t = 0, C_2 = 0 \text{ so } s = -5t^2 + 88t. \text{ At } t = 8.8, s = 387.2. \text{ The car travels 387.2 ft before coming to a stop.}$$

33.  $a = a_0 \text{ ft/s}^2$ ,  $v = a_0 t + v_0 = a_0 t + 132 \text{ ft/s}$ ,  $s = a_0 t^2/2 + 132t + s_0 = a_0 t^2/2 + 132t \text{ ft}$ ;  $s = 200 \text{ ft}$  when  $v = 88 \text{ ft/s}$ . Solve  $88 = a_0 t + 132$  and  $200 = a_0 t^2/2 + 132t$  to get  $a_0 = -\frac{121}{5}$  when  $t = \frac{20}{11}$ , so  $s = -12.1t^2 + 132t$ ,  $v = -\frac{121}{5}t + 132$ .

(a)  $a_0 = -\frac{121}{5} \text{ ft/s}^2$

(b)  $v = 55 \text{ mi/h} = \frac{242}{3} \text{ ft/s}$  when  $t = \frac{70}{33} \text{ s}$

(c)  $v = 0$  when  $t = \frac{60}{11} \text{ s}$

- 34.**  $dv/dt = 3$ ,  $v = 3t + C_1$ , but  $v = v_0$  when  $t = 0$  so  $C_1 = v_0$ ,  $v = 3t + v_0$ . From  $ds/dt = v = 3t + v_0$  we get  $s = 3t^2/2 + v_0 t + C_2$  and, with  $s = 0$  when  $t = 0$ ,  $C_2 = 0$  so  $s = 3t^2/2 + v_0 t$ .  $s = 40$  when  $t = 4$  thus  $40 = 3(4)^2/2 + v_0(4)$ ,  $v_0 = 4$  m/s
- 35.** Suppose  $s = s_0 = 0$ ,  $v = v_0 = 0$  at  $t = t_0 = 0$ ;  $s = s_1 = 120$ ,  $v = v_1$  at  $t = t_1$ ; and  $s = s_2$ ,  $v = v_2 = 12$  at  $t = t_2$ . From Exercise 30(a),  

$$2.6 = a = \frac{v_1^2 - v_0^2}{2(s_1 - s_0)}$$
,  $v_1^2 = 2as_1 = 5.2(120) = 624$ . Applying the formula again,  

$$-1.5 = a = \frac{v_2^2 - v_1^2}{2(s_2 - s_1)}$$
,  $v_2^2 = v_1^2 - 3(s_2 - s_1)$ , so  

$$s_2 = s_1 - (v_2^2 - v_1^2)/3 = 120 - (144 - 624)/3 = 280$$
 m.
- 36.**  $a(t) = \begin{cases} 4, & t < 2 \\ 0, & t > 2 \end{cases}$ , so, with  $v_0 = 0$ ,  $v(t) = \begin{cases} 4t, & t < 2 \\ 8, & t > 2 \end{cases}$  and,  
since  $s_0 = 0$ ,  $s(t) = \begin{cases} 2t^2, & t < 2 \\ 8t - 8, & t > 2 \end{cases}$   $s = 100$  when  $8t - 8 = 100$ ,  $t = 108/8 = 13.5$  s
- 37.** The truck's velocity is  $v_T = 50$  and its position is  $s_T = 50t + 5000$ . The car's acceleration is  $a_C = 2$ , so  $v_C = 2t$ ,  $s_C = t^2$  (initial position and initial velocity of the car are both zero).  $s_T = s_C$  when  $50t + 5000 = t^2$ ,  $t^2 - 50t - 5000 = (t+50)(t-100) = 0$ ,  $t = 100$  s and  $s_C = s_T = t^2 = 10,000$  ft.
- 38.** Let  $t = 0$  correspond to the time when the leader is 100 m from the finish line; let  $s = 0$  correspond to the finish line. Then  $v_C = 12$ ,  $s_C = 12t - 115$ ;  $a_L = 0.5$  for  $t > 0$ ,  $v_L = 0.5t + 8$ ,  $s_L = 0.25t^2 + 8t - 100$ .  $s_C = 0$  at  $t = 115/12 \approx 9.58$  s, and  $s_L = 0$  at  $t = -16 + 4\sqrt{41} \approx 9.61$ , so the challenger wins.
- 39.**  $s = 0$  and  $v = 112$  when  $t = 0$  so  $v(t) = -32t + 112$ ,  $s(t) = -16t^2 + 112t$ 
  - (a)  $v(3) = 16$  ft/s,  $v(5) = -48$  ft/s
  - (b)  $v = 0$  when the projectile is at its maximum height so  $-32t + 112 = 0$ ,  $t = 7/2$  s,  $s(7/2) = -16(7/2)^2 + 112(7/2) = 196$  ft.
  - (c)  $s = 0$  when it reaches the ground so  $-16t^2 + 112t = 0$ ,  $-16t(t - 7) = 0$ ,  $t = 0, 7$  of which  $t = 7$  is when it is at ground level on its way down.  $v(7) = -112$ ,  $|v| = 112$  ft/s.
- 40.**  $s = 112$  when  $t = 0$  so  $s(t) = -16t^2 + v_0t + 112$ . But  $s = 0$  when  $t = 2$  thus  $-16(2)^2 + v_0(2) + 112 = 0$ ,  $v_0 = -24$  ft/s.
- 41.** (a)  $s(t) = 0$  when it hits the ground,  $s(t) = -16t^2 + 16t = -16t(t - 1) = 0$  when  $t = 1$  s.  
(b) The projectile moves upward until it gets to its highest point where  $v(t) = 0$ ,  $v(t) = -32t + 16 = 0$  when  $t = 1/2$  s.
- 42.** (a)  $s(t) = 0$  when the rock hits the ground,  $s(t) = -16t^2 + 555 = 0$  when  $t = \sqrt{555}/4$  s  
(b)  $v(t) = -32t$ ,  $v(\sqrt{555}/4) = -8\sqrt{555}$ , the speed at impact is  $8\sqrt{555}$  ft/s
- 43.** (a)  $s(t) = 0$  when the package hits the ground,  
 $s(t) = -16t^2 + 20t + 200 = 0$  when  $t = (5 + 5\sqrt{33})/8$  s  
(b)  $v(t) = -32t + 20$ ,  $v[(5 + 5\sqrt{33})/8] = -20\sqrt{33}$ , the speed at impact is  $20\sqrt{33}$  ft/s
- 44.** (a)  $s(t) = 0$  when the stone hits the ground,  
 $s(t) = -16t^2 - 96t + 112 = -16(t^2 + 6t - 7) = -16(t + 7)(t - 1) = 0$  when  $t = 1$  s  
(b)  $v(t) = -32t - 96$ ,  $v(1) = -128$ , the speed at impact is 128 ft/s

45.  $s(t) = -4.9t^2 + 49t + 150$  and  $v(t) = -9.8t + 49$
- the projectile reaches its maximum height when  $v(t) = 0$ ,  $-9.8t + 49 = 0$ ,  $t = 5$  s
  - $s(5) = -4.9(5)^2 + 49(5) + 150 = 272.5$  m
  - the projectile reaches its starting point when  $s(t) = 150$ ,  $-4.9t^2 + 49t + 150 = 150$ ,  $-4.9t(t - 10) = 0$ ,  $t = 10$  s
  - $v(10) = -9.8(10) + 49 = -49$  m/s
  - $s(t) = 0$  when the projectile hits the ground,  $-4.9t^2 + 49t + 150 = 0$  when (use the quadratic formula)  $t \approx 12.46$  s
  - $v(12.46) = -9.8(12.46) + 49 \approx -73.1$ , the speed at impact is about 73.1 m/s
46. take  $s = 0$  at the water level and let  $h$  be the height of the bridge, then  $s = h$  and  $v = 0$  when  $t = 0$  so  $s(t) = -16t^2 + h$
- $s = 0$  when  $t = 4$  thus  $-16(4)^2 + h = 0$ ,  $h = 256$  ft
  - First, find how long it takes for the stone to hit the water (find  $t$  for  $s = 0$ ) :  $-16t^2 + h = 0$ ,  $t = \sqrt{h}/4$ . Next, find how long it takes the sound to travel to the bridge: this time is  $h/1080$  because the speed is constant at 1080 ft/s. Finally, use the fact that the total of these two times must be 4 s:  $\frac{h}{1080} + \frac{\sqrt{h}}{4} = 4$ ,  $h + 270\sqrt{h} = 4320$ ,  $h + 270\sqrt{h} - 4320 = 0$ , and by the quadratic formula  $\sqrt{h} = \frac{-270 \pm \sqrt{(270)^2 + 4(4320)}}{2}$ , reject the negative value to get  $\sqrt{h} \approx 15.15$ ,  $h \approx 229.5$  ft.
47.  $g = 9.8/6 = 4.9/3$  m/s<sup>2</sup>, so  $v = -(4.9/3)t$ ,  $s = -(4.9/6)t^2 + 5$ ,  $s = 0$  when  $t = \sqrt{30/4.9}$  and  $v = -(4.9/3)\sqrt{30/4.9} \approx -4.04$ , so the speed of the module upon landing is 4.04 m/s
48.  $s(t) = -\frac{1}{2}gt^2 + v_0t$ ;  $s = 1000$  when  $v = 0$ , so  $0 = v = -gt + v_0$ ,  $t = v_0/g$ ,  $1000 = s(v_0/g) = -\frac{1}{2}g(v_0/g)^2 + v_0(v_0/g) = \frac{1}{2}v_0^2/g$ , so  $v_0^2 = 2000g$ ,  $v_0 = \sqrt{2000g}$ . The initial velocity on the Earth would have to be  $\sqrt{6}$  times faster than that on the Moon.

49.  $f_{\text{ave}} = \frac{1}{3-1} \int_1^3 3x \, dx = \left[ \frac{3}{4}x^2 \right]_1^3 = 6$


50.  $f_{\text{ave}} = \frac{1}{2-(-1)} \int_{-1}^2 x^2 \, dx = \left[ \frac{1}{9}x^3 \right]_{-1}^2 = 1$

51.  $f_{\text{ave}} = \frac{1}{\pi-0} \int_0^\pi \sin x \, dx = \left[ -\frac{1}{\pi} \cos x \right]_0^\pi = 2/\pi$


52.  $f_{\text{ave}} = \frac{1}{\pi-0} \int_0^\pi \cos x \, dx = \left[ \frac{1}{\pi} \sin x \right]_0^\pi = 0$

53. (a)  $f_{\text{ave}} = \frac{1}{2-0} \int_0^2 x^2 \, dx = 4/3$

(b)  $(x^*)^2 = 4/3$ ,  $x^* = \pm 2/\sqrt{3}$ , but only  $2/\sqrt{3}$  is in  $[0, 2]$



54. (a)  $f_{\text{ave}} = \frac{1}{4-0} \int_0^4 2x \, dx = 4$  (b)  $2x^* = 4, x^* = 2$



55. (a)  $v_{\text{ave}} = \frac{1}{4-1} \int_1^4 (3t^3 + 2) \, dt = \frac{1}{3} \cdot \frac{789}{4} = \frac{263}{4}$

(b)  $v_{\text{ave}} = \frac{s(4) - s(1)}{4-1} = \frac{100 - 7}{3} = 31$

56. (a)  $a_{\text{ave}} = \frac{1}{5-0} \int_0^5 (t+1) \, dt = 7/2$

(b)  $a_{\text{ave}} = \frac{v(\pi/4) - v(0)}{\pi/4 - 0} = \frac{\sqrt{2}/2 - 1}{\pi/4} = (2\sqrt{2} - 4)/\pi$

57. time to fill tank = (volume of tank)/(rate of filling) =  $[\pi(3)^2 5]/(1) = 45\pi$ , weight of water in tank at time  $t = (62.4)$  (rate of filling)(time) =  $62.4t$ ,

$$\text{weight}_{\text{ave}} = \frac{1}{45\pi} \int_0^{45\pi} 62.4t \, dt = 1404\pi \text{ lb}$$

58. (a) If  $x$  is the distance from the cooler end, then the temperature is  $T(x) = (15 + 1.5x)^\circ \text{ C}$ , and  $T_{\text{ave}} = \frac{1}{10-0} \int_0^{10} (15 + 1.5x) \, dx = 22.5^\circ \text{ C}$

- (b) By the Mean-Value Theorem for Integrals there exists  $x^*$  in  $[0, 10]$  such that  $f(x^*) = \frac{1}{10-0} \int_0^{10} (15 + 1.5x) \, dx = 22.5, 15 + 1.5x^* = 22.5, x^* = 5$

59. (a) amount of water = (rate of flow)(time) =  $4t$  gal, total amount =  $4(30) = 120$  gal

(b) amount of water =  $\int_0^{60} (4 + t/10) \, dt = 420$  gal

(c) amount of water =  $\int_0^{120} (10 + \sqrt{t}) \, dt = 1200 + 160\sqrt{30} \approx 2076.36$  gal

60. (a) The maximum value of  $R$  occurs at 4:30 P.M. when  $t = 0$ .

(b)  $\int_0^{60} 100(1 - 0.0001t^2) \, dt = 5280$  cars

61. (a)  $\int_a^b [f(x) - f_{\text{ave}}] \, dx = \int_a^b f(x) \, dx - \int_a^b f_{\text{ave}} \, dx = \int_a^b f(x) \, dx - f_{\text{ave}}(b-a) = 0$

because  $f_{\text{ave}}(b-a) = \int_a^b f(x) \, dx$

(b) no, because if  $\int_a^b [f(x) - c]dx = 0$  then  $\int_a^b f(x)dx - c(b-a) = 0$  so  
 $c = \frac{1}{b-a} \int_a^b f(x)dx = f_{\text{ave}}$  is the only value

## EXERCISE SET 5.8

1. (a)  $\int_1^3 u^7 du$       (b)  $-\frac{1}{2} \int_7^4 u^{1/2} du$       (c)  $\frac{1}{\pi} \int_{-\pi}^{\pi} \sin u du$       (d)  $\int_{-3}^0 (u+5)u^{20} du$

2. (a)  $\frac{1}{2} \int_{-3}^7 u^8 du$       (b)  $\int_{3/2}^{5/2} \frac{1}{\sqrt{u}} du$

(c)  $\int_0^1 u^2 du$       (d)  $\frac{1}{2} \int_3^4 (u-3)u^{1/2} du$

3.  $u = 2x + 1$ ,  $\frac{1}{2} \int_1^3 u^4 du = \frac{1}{10} u^5 \Big|_1^3 = 121/5$ , or  $\frac{1}{10} (2x+1)^5 \Big|_0^1 = 121/5$

4.  $u = 4x - 2$ ,  $\frac{1}{4} \int_2^6 u^3 du = \frac{1}{16} u^4 \Big|_2^6 = 80$ , or  $\frac{1}{16} (4x-2)^4 \Big|_1^2 = 80$

5.  $u = 1 - 2x$ ,  $-\frac{1}{2} \int_3^1 u^3 du = -\frac{1}{8} u^4 \Big|_3^1 = 10$ , or  $-\frac{1}{8} (1-2x)^4 \Big|_{-1}^0 = 10$

6.  $u = 4 - 3x$ ,  $-\frac{1}{3} \int_1^{-2} u^8 du = -\frac{1}{27} u^9 \Big|_1^{-2} = 19$ , or  $-\frac{1}{27} (4-3x)^9 \Big|_1^{-2} = 19$

7.  $u = 1 + x$ ,  $\int_1^9 (u-1)u^{1/2} du = \int_1^9 (u^{3/2} - u^{1/2}) du = \frac{2}{5} u^{5/2} - \frac{2}{3} u^{3/2} \Big|_1^9 = 1192/15$ ,  
or  $\frac{2}{5} (1+x)^{5/2} - \frac{2}{3} (1+x)^{3/2} \Big|_0^8 = 1192/15$

8.  $u = 4 - x$ ,  $\int_9^4 (u-4)u^{1/2} du = \int_9^4 (u^{3/2} - 4u^{1/2}) du = \frac{2}{5} u^{5/2} - \frac{8}{3} u^{3/2} \Big|_9^4 = -506/15$   
or  $\frac{2}{5} (4-x)^{5/2} - \frac{8}{3} (4-x)^{3/2} \Big|_{-5}^0 = -506/15$

9.  $u = x/2$ ,  $8 \int_0^{\pi/4} \sin u du = -8 \cos u \Big|_0^{\pi/4} = 8 - 4\sqrt{2}$ , or  $-8 \cos(x/2) \Big|_0^{\pi/2} = 8 - 4\sqrt{2}$

10.  $u = 3x$ ,  $\frac{2}{3} \int_0^{\pi/2} \cos u du = \frac{2}{3} \sin u \Big|_0^{\pi/2} = 2/3$ , or  $\frac{2}{3} \sin 3x \Big|_0^{\pi/6} = 2/3$

11.  $u = x^2 + 2$ ,  $\frac{1}{2} \int_6^3 u^{-3} du = -\frac{1}{4u^2} \Big|_6^3 = -1/48$ , or  $-\frac{1}{4} \frac{1}{(x^2+2)^2} \Big|_{-2}^{-1} = -1/48$

**12.**  $u = \frac{1}{4}x - \frac{1}{4}$ ,  $4 \int_{-\pi/4}^{\pi/4} \sec^2 u du = 4 \tan u \Big|_{-\pi/4}^{\pi/4} = 8$ , or  $4 \tan \left( \frac{1}{4}x - \frac{1}{4} \right) \Big|_{1-\pi}^{1+\pi} = 8$

**13.**  $\frac{1}{3} \int_0^5 \sqrt{25-u^2} du = \frac{1}{3} \left[ \frac{1}{4}\pi(5)^2 \right] = \frac{25}{12}\pi$       **14.**  $\frac{1}{2} \int_0^4 \sqrt{16-u^2} du = \frac{1}{2} \left[ \frac{1}{4}\pi(4)^2 \right] = 2\pi$

**15.**  $-\frac{1}{2} \int_1^0 \sqrt{1-u^2} du = \frac{1}{2} \int_0^1 \sqrt{1-u^2} du = \frac{1}{2} \cdot \frac{1}{4}[\pi(1)^2] = \pi/8$

**16.**  $\int_{-2}^2 \sqrt{4-u^2} du = \frac{1}{2}[\pi(2)^2] = 2\pi$

**17.**  $\int_0^1 \sin \pi x dx = -\frac{1}{\pi} \cos \pi x \Big|_0^1 = -\frac{1}{\pi}(-1-1) = 2/\pi$

**18.**  $A = \int_0^{\pi/8} 3 \cos 2x dx = \frac{3}{2} \sin 2x \Big|_0^{\pi/8} = 3\sqrt{2}/4$

**19.**  $\int_3^7 (x+5)^{-2} dx = -(x+5)^{-1} \Big|_3^7 = -\frac{1}{12} + \frac{1}{8} = \frac{1}{24}$

**20.**  $A = \int_0^1 \frac{dx}{(3x+1)^2} = -\frac{1}{3(3x+1)} \Big|_0^1 = \frac{1}{4}$

**21.**  $\frac{1}{2-0} \int_0^2 \frac{x}{(5x^2+1)^2} dx = -\frac{1}{2} \frac{1}{10} \frac{1}{5x^2+1} \Big|_0^2 = \frac{1}{21}$

**22.**  $f_{\text{ave}} = \frac{1}{1/4 - (-1/4)} \int_{-1/4}^{1/4} \sec^2 \pi x dx = \frac{2}{\pi} \tan \pi x \Big|_{-1/4}^{1/4} = \frac{4}{\pi}$

**23.**  $\frac{2}{3}(3x+1)^{1/2} \Big|_0^1 = 2/3$

**24.**  $\frac{2}{15}(5x-1)^{3/2} \Big|_1^2 = 38/15$

**25.**  $\frac{2}{3}(x^3+9)^{1/2} \Big|_{-1}^1 = \frac{2}{3}(\sqrt{10}-2\sqrt{2})$

**26.**  $\frac{1}{10}(t^3+1)^{20} \Big|_{-1}^0 = 1/10$

**27.**  $u = x^2 + 4x + 7$ ,  $\frac{1}{2} \int_{12}^{28} u^{-1/2} du = u^{1/2} \Big|_{12}^{28} = \sqrt{28} - \sqrt{12} = 2(\sqrt{7} - \sqrt{3})$

**28.**  $\int_1^2 \frac{1}{(x-3)^2} dx = -\frac{1}{x-3} \Big|_1^2 = 1/2$

**29.**  $\frac{1}{2} \sin^2 x \Big|_{-3\pi/4}^{\pi/4} = 0$

**30.**  $\frac{2}{3}(\tan x)^{3/2} \Big|_0^{\pi/4} = 2/3$

**31.**  $\frac{5}{2} \sin(x^2) \Big|_0^{\sqrt{\pi}} = 0$

**32.**  $u = \sqrt{x}$ ,  $2 \int_{\pi}^{2\pi} \sin u du = -2 \cos u \Big|_{\pi}^{2\pi} = -4$

33.  $u = 3\theta, \frac{1}{3} \int_{\pi/4}^{\pi/3} \sec^2 u du = \frac{1}{3} \tan u \Big|_{\pi/4}^{\pi/3} = (\sqrt{3} - 1)/3$

34.  $u = \sin 3\theta, \frac{1}{3} \int_0^{-1} u^2 du = \frac{1}{9} u^3 \Big|_0^{-1} = -1/9$

35.  $u = 4 - 3y, y = \frac{1}{3}(4 - u), dy = -\frac{1}{3}du$   
 $-\frac{1}{27} \int_4^1 \frac{16 - 8u + u^2}{u^{1/2}} du = \frac{1}{27} \int_1^4 (16u^{-1/2} - 8u^{1/2} + u^{3/2}) du$   
 $= \frac{1}{27} \left[ 32u^{1/2} - \frac{16}{3}u^{3/2} + \frac{2}{5}u^{5/2} \right]_1^4 = 106/405$

36.  $u = 5 + x, \int_4^9 \frac{u-5}{\sqrt{u}} du = \int_4^9 (u^{1/2} - 5u^{-1/2}) du = \frac{2}{3}u^{3/2} - 10u^{1/2} \Big|_4^9 = 8/3$

37. (b)  $\int_0^{\pi/6} \sin^4 x (1 - \sin^2 x) \cos x dx = \left( \frac{1}{5} \sin^5 x - \frac{1}{7} \sin^7 x \right) \Big|_0^{\pi/6} = \frac{1}{160} - \frac{1}{896} = \frac{23}{4480}$

38. (b)  $\int_{-\pi/4}^{\pi/4} \tan^2 x (\sec^2 x - 1) dx = \frac{1}{3} \tan^3 x \Big|_{-\pi/4}^{\pi/4} - \int_{-\pi/4}^{\pi/4} (\sec^2 x - 1) dx$   
 $= \frac{2}{3} + (-\tan x + x) \Big|_{-\pi/4}^{\pi/4} = \frac{2}{3} - 2 + \frac{\pi}{2} = -\frac{4}{3} + \frac{\pi}{2}$

39. (a)  $u = 3x + 1, \frac{1}{3} \int_1^4 f(u) du = 5/3$       (b)  $u = 3x, \frac{1}{3} \int_0^9 f(u) du = 5/3$

(c)  $u = x^2, 1/2 \int_4^0 f(u) du = -1/2 \int_0^4 f(u) du = -1/2$

40.  $u = 1 - x, \int_0^1 x^m (1 - x)^n dx = - \int_1^0 (1 - u)^m u^n du = \int_0^1 u^n (1 - u)^m du = \int_0^1 x^n (1 - x)^m dx$

41.  $\sin x = \cos(\pi/2 - x),$   
 $\int_0^{\pi/2} \sin^n x dx = \int_0^{\pi/2} \cos^n(\pi/2 - x) dx = - \int_{\pi/2}^0 \cos^n u du \quad (u = \pi/2 - x)$   
 $= \int_0^{\pi/2} \cos^n u du = \int_0^{\pi/2} \cos^n x dx \quad (\text{by replacing } u \text{ by } x)$

42.  $u = 1 - x, - \int_1^0 (1 - u) u^n du = \int_0^1 (1 - u) u^n du = \int_0^1 (u^n - u^{n+1}) du = \frac{1}{n+1} - \frac{1}{n+2}$   
 $= \frac{1}{(n+1)(n+2)}$

43. (a)  $V_{\text{rms}}^2 = \frac{1}{1/f - 0} \int_0^{1/f} V_p^2 \sin^2(2\pi ft) dt = \frac{1}{2} f V_p^2 \int_0^{1/f} [1 - \cos(4\pi ft)] dt$   
 $= \frac{1}{2} f V_p^2 \left[ t - \frac{1}{4\pi f} \sin(4\pi ft) \right]_0^{1/f} = \frac{1}{2} V_p^2, \text{ so } V_{\text{rms}} = V_p/\sqrt{2}$

(b)  $V_p/\sqrt{2} = 120, V_p = 120\sqrt{2} \approx 169.7 \text{ V}$

44. Let  $u = t - x$ , then  $du = -dx$  and

$$\int_0^t f(t-x)g(x)dx = - \int_t^0 f(u)g(t-u)du = \int_0^t f(u)g(t-u)du;$$

the result follows by replacing  $u$  by  $x$  in the last integral.

45. (a)  $I = - \int_a^0 \frac{f(a-u)}{f(a-u) + f(u)} du = \int_0^a \frac{f(a-u) + f(u) - f(u)}{f(a-u) + f(u)} du$   
 $= \int_0^a du - \int_0^a \frac{f(u)}{f(a-u) + f(u)} du, I = a - I \text{ so } 2I = a, I = a/2$

(b) 3/2

(c)  $\pi/4$

46.  $x = \frac{1}{u}, dx = -\frac{1}{u^2}du, I = \int_{-1}^1 \frac{1}{1+1/u^2}(-1/u^2)du = - \int_{-1}^1 \frac{1}{u^2+1}du = -I$  so  $I = 0$  which is impossible because  $\frac{1}{1+x^2}$  is positive on  $[-1, 1]$ . The substitution  $u = 1/x$  is not valid because  $u$  is not continuous for all  $x$  in  $[-1, 1]$ .

47.  $\int_0^1 \sin \pi x dx = 2/\pi$

49. (a) Let  $u = -x$  then

$$\int_{-a}^a f(x)dx = - \int_a^{-a} f(-u)du = \int_{-a}^a f(-u)du = - \int_{-a}^a f(u)du$$

so, replacing  $u$  by  $x$  in the latter integral,

$$\int_{-a}^a f(x)dx = - \int_{-a}^a f(x)dx, 2 \int_{-a}^a f(x)dx = 0, \int_{-a}^a f(x)dx = 0$$

The graph of  $f$  is symmetric about the origin so  $\int_{-a}^0 f(x)dx$  is the negative of  $\int_0^a f(x)dx$   
thus  $\int_{-a}^a f(x)dx = \int_{-a}^0 f(x)dx + \int_0^a f(x)dx = 0$

(b)  $\int_{-a}^a f(x)dx = \int_{-a}^0 f(x)dx + \int_0^a f(x)dx$ , let  $u = -x$  in  $\int_{-a}^0 f(x)dx$  to get

$$\int_{-a}^0 f(x)dx = - \int_a^0 f(-u)du = \int_0^a f(-u)du = \int_0^a f(u)du = \int_0^a f(x)dx$$

so  $\int_{-a}^a f(x)dx = \int_0^a f(x)dx + \int_0^a f(x)dx = 2 \int_0^a f(x)dx$

The graph of  $f(x)$  is symmetric about the  $y$ -axis so there is as much signed area to the left of the  $y$ -axis as there is to the right.

50. (a) By Exercise 49(a),  $\int_{-1}^1 x \sqrt{\cos(x^2)} dx = 0$

(b)  $u = x - \pi/2, du = dx, \sin(u + \pi/2) = \sin u, \cos(u + \pi/2) = -\sin u$

$$\int_0^\pi \sin^8 x \cos^5 x dx = \int_{-\pi/2}^{\pi/2} \sin^8 u (-\sin^5 u) du = - \int_{-\pi/2}^{\pi/2} \sin^{13} u du = 0 \text{ by Exercise 49(a).}$$

## CHAPTER 5 SUPPLEMENTARY EXERCISES

5. If the acceleration  $a = \text{const}$ , then  $v(t) = at + v_0, s(t) = \frac{1}{2}at^2 + v_0t + s_0$ .

6. (a) Divide the base into  $n$  equal subintervals. Above each subinterval choose the lowest and highest points on the curved top. Draw a rectangle above the subinterval going through the lowest point, and another through the highest point. Add the rectangles that go through the lowest points to obtain a lower estimate of the area; add the rectangles through the highest points to obtain an upper estimate of the area.

(b)  $n = 10$ : 25.0 cm, 22.4 cm

(c)  $n = 20$ : 24.4 cm, 23.1 cm

7. (a)  $\frac{1}{2} + \frac{1}{4} = \frac{3}{4}$

(b)  $-1 - \frac{1}{2} = -\frac{3}{2}$

(c)  $5 \left( -1 - \frac{3}{4} \right) = -\frac{35}{4}$

(d) -2

(e) not enough information

(f) not enough information

8. (a)  $\frac{1}{2} + 2 = \frac{5}{2}$

(b) not enough information

(c) not enough information


(d)  $4(2) - 3\frac{1}{2} = \frac{13}{2}$

9. (a)  $\int_{-1}^1 dx + \int_{-1}^1 \sqrt{1-x^2} dx = 2(1) + \pi(1)^2/2 = 2 + \pi/2$

(b)  $\frac{1}{3}(x^2+1)^{3/2} \Big|_0^3 - \pi(3)^2/4 = \frac{1}{3}(10^{3/2} - 1) - 9\pi/4$

(c)  $u = x^2, du = 2xdx; \frac{1}{2} \int_0^1 \sqrt{1-u^2} du = \frac{1}{2}\pi(1)^2/4 = \pi/8$

10.  $\frac{1}{2}$



11. The rectangle with vertices  $(0,0)$ ,  $(\pi,0)$ ,  $(\pi,1)$  and  $(0,1)$  has area  $\pi$  and is much too large; so is the triangle with vertices  $(0,0)$ ,  $(\pi,0)$  and  $(\pi,1)$  which has area  $\pi/2$ ;  $1 - \pi$  is negative; so the answer is  $35\pi/128$ .

12. (a)  $\frac{1}{n} \sum_{k=1}^n \sqrt{\frac{k}{n}} = \sum_{k=1}^n f(x_k^*) \Delta x$  where  $f(x) = \sqrt{x}$ ,  $x_k^* = k/n$ , and  $\Delta x = 1/n$  for  $0 \leq x \leq 1$ . Thus

$$\lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{k=1}^n \sqrt{\frac{k}{n}} = \int_0^1 x^{1/2} dx = \frac{2}{3}$$

(b)  $\frac{1}{n} \sum_{k=1}^n \left(\frac{k}{n}\right)^4 = \sum_{k=1}^n f(x_k^*) \Delta x$  where  $f(x) = x^4$ ,  $x_k^* = k/n$ , and  $\Delta x = 1/n$  for  $0 \leq x \leq 1$ . Thus

$$\lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{k=1}^n \left(\frac{k}{n}\right)^4 = \int_0^1 x^4 dx = \frac{1}{5}$$

13. left endpoints:  $x_k^* = 1, 2, 3, 4$ ;  $\sum_{k=1}^4 f(x_k^*) \Delta x = (2+3+2+1)(1) = 8$

right endpoints:  $x_k^* = 2, 3, 4, 5$ ;  $\sum_{k=1}^4 f(x_k^*) \Delta x = (3+2+1+2)(1) = 8$

14. The direction field is clearly an even function, which means that the solution is even, its derivative is odd. Since  $\sin x$  is periodic and the direction field is not, that eliminates all but  $x$ , the solution of which is the family  $y = x^2/2 + C$ .

15. (a)  $1 \cdot 2 + 2 \cdot 3 + \cdots + n(n+1) = \sum_{k=1}^n k(k+1) = \sum_{k=1}^n k^2 + \sum_{k=1}^n k$   
 $= \frac{1}{6}n(n+1)(2n+1) + \frac{1}{2}n(n+1) = \frac{1}{3}n(n+1)(n+2)$

(b)  $\sum_{k=1}^{n-1} \left( \frac{9}{n} - \frac{k}{n^2} \right) = \frac{9}{n} \sum_{k=1}^{n-1} 1 - \frac{1}{n^2} \sum_{k=1}^{n-1} k = \frac{9}{n}(n-1) - \frac{1}{n^2} \cdot \frac{1}{2}(n-1)(n) = \frac{17}{2} \left( \frac{n-1}{n} \right);$

$$\lim_{n \rightarrow +\infty} \frac{17}{2} \left( \frac{n-1}{n} \right) = \frac{17}{2}$$

(c)  $\sum_{i=1}^3 \left[ \sum_{j=1}^2 i + \sum_{j=1}^2 j \right] = \sum_{i=1}^3 \left[ 2i + \frac{1}{2}(2)(3) \right] = 2 \sum_{i=1}^3 i + \sum_{i=1}^3 3 = 2 \cdot \frac{1}{2}(3)(4) + (3)(3) = 21$

16. (a)  $\sum_{k=0}^{14} (k+4)(k+1)$  (b)  $\sum_{k=5}^{19} (k-1)(k-4)$

17. For  $1 \leq k \leq n$  the  $k$ -th  $L$ -shaped strip consists of the corner square, a strip above and a strip to the right for a combined area of  $1 + (k-1) + (k-1) = 2k-1$ , so the total area is  $\sum_{k=1}^n (2k-1) = n^2$ .

**18.**  $1 + 3 + 5 + \cdots + (2n - 1) = \sum_{k=1}^n (2k - 1) = 2 \sum_{k=1}^n k - \sum_{k=1}^n 1 = 2 \cdot \frac{1}{2}n(n+1) - n = n^2$

**19.**  $(3^5 - 3^4) + (3^6 - 3^5) + \cdots + (3^{17} - 3^{16}) = 3^{17} - 3^4$

**20.**  $\left(1 - \frac{1}{2}\right) + \left(\frac{1}{2} - \frac{1}{3}\right) + \cdots + \left(\frac{1}{50} - \frac{1}{51}\right) = \frac{50}{51}$

**21.**  $\left(\frac{1}{2^2} - \frac{1}{1^2}\right) + \left(\frac{1}{3^2} - \frac{1}{2^2}\right) + \cdots + \left(\frac{1}{20^2} - \frac{1}{19^2}\right) = \frac{1}{20^2} - 1 = -\frac{399}{400}$

**22.**  $(2^2 - 2) + (2^3 - 2^2) + \cdots + (2^{101} - 2^{100}) = 2^{101} - 2$

**23. (a)** 
$$\begin{aligned} \sum_{k=1}^n \frac{1}{(2k-1)(2k+1)} &= \frac{1}{2} \sum_{k=1}^n \left( \frac{1}{2k-1} - \frac{1}{2k+1} \right) \\ &= \frac{1}{2} \left[ \left(1 - \frac{1}{3}\right) + \left(\frac{1}{3} - \frac{1}{5}\right) + \left(\frac{1}{5} - \frac{1}{7}\right) + \cdots + \left(\frac{1}{2n-1} - \frac{1}{2n+1}\right) \right] \\ &= \frac{1}{2} \left[ 1 - \frac{1}{2n+1} \right] = \frac{n}{2n+1} \end{aligned}$$

**(b)**  $\lim_{n \rightarrow +\infty} \frac{n}{2n+1} = \frac{1}{2}$

**24. (a)** 
$$\begin{aligned} \sum_{k=1}^n \frac{1}{k(k+1)} &= \sum_{k=1}^n \left( \frac{1}{k} - \frac{1}{k+1} \right) \\ &= \left(1 - \frac{1}{2}\right) + \left(\frac{1}{2} - \frac{1}{3}\right) + \left(\frac{1}{3} - \frac{1}{4}\right) + \cdots + \left(\frac{1}{n} - \frac{1}{n+1}\right) \\ &= 1 - \frac{1}{n+1} = \frac{n}{n+1} \end{aligned}$$

**(b)**  $\lim_{n \rightarrow +\infty} \frac{n}{n+1} = 1$

**25.**  $\sum_{i=1}^n (x_i - \bar{x}) = \sum_{i=1}^n x_i - \sum_{i=1}^n \bar{x} = \sum_{i=1}^n x_i - n\bar{x}$  but  $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$  thus

$$\sum_{i=1}^n x_i = n\bar{x} \text{ so } \sum_{i=1}^n (x_i - \bar{x}) = n\bar{x} - n\bar{x} = 0$$

**26.** 
$$\begin{aligned} S - rS &= \sum_{k=0}^n ar^k - \sum_{k=0}^n ar^{k+1} \\ &= (a + ar + ar^2 + \cdots + ar^n) - (ar + ar^2 + ar^3 + \cdots + ar^{n+1}) \\ &= a - ar^{n+1} = a(1 - r^{n+1}) \end{aligned}$$

so  $(1 - r)S = a(1 - r^{n+1})$ , hence  $S = a(1 - r^{n+1})/(1 - r)$

$$27. \quad (a) \quad \sum_{k=0}^{19} 3^{k+1} = \sum_{k=0}^{19} 3(3^k) = \frac{3(1 - 3^{20})}{1 - 3} = \frac{3}{2}(3^{20} - 1)$$

$$(b) \quad \sum_{k=0}^{25} 2^{k+5} = \sum_{k=0}^{25} 2^5 2^k = \frac{2^5(1 - 2^{26})}{1 - 2} = 2^{31} - 2^5$$

$$(c) \quad \sum_{k=0}^{100} (-1) \left( \frac{-1}{2} \right)^k = \frac{(-1)(1 - (-1/2)^{101})}{1 - (-1/2)} = -\frac{2}{3}(1 + 1/2^{101})$$

28. (a) 1.999023438, 1.999999046, 2.000000000; 2  
 (b) 2.831059456, 2.990486364, 2.999998301; 3

29. (a) If  $u = \sec x$ ,  $du = \sec x \tan x dx$ ,  $\int \sec^2 x \tan x dx = \int u du = u^2/2 + C_1 = (\sec^2 x)/2 + C_1$ ;  
 if  $u = \tan x$ ,  $du = \sec^2 x dx$ ,  $\int \sec^2 x \tan x dx = \int u du = u^2/2 + C_2 = (\tan^2 x)/2 + C_2$ .

- (b) They are equal only if  $\sec^2 x$  and  $\tan^2 x$  differ by a constant, which is true.

**30.**  $\frac{1}{2} \sec^2 x \Big|_0^{\pi/4} = \frac{1}{2}(2 - 1) = 1/2$  and  $\frac{1}{2} \tan^2 x \Big|_0^{\pi/4} = \frac{1}{2}(1 - 0) = 1/2$

$$31. \quad \int \sqrt{1+x^{-2/3}} dx = \int x^{-1/3} \sqrt{x^{2/3}+1} dx; \quad u = x^{2/3} + 1, \quad du = \frac{2}{3}x^{-1/3}dx$$

$$\frac{3}{2} \int u^{1/2} du = u^{3/2} + C = (x^{2/3} + 1)^{3/2} + C$$

$$32. \quad (a) \quad \int_a^b \sum_{k=1}^n f_k(x) dx = \sum_{k=1}^n \int_a^b f_k(x) dx$$

- (b) yes; substitute  $c_k f_k(x)$  for  $f_k(x)$  in part (a), and then use  $\int_a^b c_k f_k(x) dx = c_k \int_a^b f_k(x) dx$  from Theorem 5.5.4


33. (a)  $\int_1^x \frac{1}{1+t^2} dt$

$$(b) \quad \int_{\tan(\pi/4-2)}^x \frac{1}{1+t^2} dt$$

**34. (a)**  $F'(x) = \frac{x-3}{x^2+7}$ ; increasing on  $[3, +\infty)$ , decreasing on  $(-\infty, 3]$

(b)  $F''(x) = \frac{7+6x-x^2}{(x^2+7)^2} = \frac{(7-x)(1+x)}{(x^2+7)^2}$ ; concave up on  $(-1, 7)$ , concave down on  $(-\infty, -1)$  and  $(7, +\infty)$

- (c)  $F'(x) = \frac{x-3}{x^2+7} = 0$  when  $x = 3$ , which is a relative minimum, and hence the absolute minimum, by the first derivative test.



**35.**  $F'(x) = \frac{1}{1+x^2} + \frac{1}{1+(1/x)^2}(-1/x^2) = 0$  so  $F$  is constant on  $(0, +\infty)$ .

- 36.**  $(-3, 3)$  because  $f$  is continuous there and 1 is in  $(-3, 3)$

- 37.** (a) The domain is  $(-\infty, +\infty)$ ;  $F(x)$  is 0 if  $x = 1$ , positive if  $x > 1$ , and negative if  $x < 1$ , because the integrand is positive, so the sign of the integral depends on the orientation (forwards or backwards).  
 (b) The domain is  $[-2, 2]$ ;  $F(x)$  is 0 if  $x = -1$ , positive if  $-1 < x \leq 2$ , and negative if  $-2 \leq x < -1$ ; same reasons as in Part (a).

**38.** The left endpoint of the top boundary is  $((b-a)/2, h)$  and the right endpoint of the top boundary is  $((b+a)/2, h)$  so

$$f(x) = \begin{cases} 2hx/(b-a), & x < (b-a)/2 \\ h, & (b-a)/2 < x < (b+a)/2 \\ 2h(x-b)/(a-b), & x > (a+b)/2 \end{cases}$$

The area of the trapezoid is given by

$$\int_0^{(b-a)/2} \frac{2hx}{b-a} dx + \int_{(b-a)/2}^{(b+a)/2} hdx + \int_{(b+a)/2}^b \frac{2h(x-b)}{a-b} dx = (b-a)h/4 + ah + (b-a)h/4 = h(a+b)/2.$$



**40.**  $w(t) = \int_0^t \tau/7 d\tau = t^2/14$ , assuming  $w_0 = w(0) = 0$ ;  $w_{\text{ave}} = \frac{1}{26} \int_{26}^{52} t^2/7 dt = \frac{1}{26} \frac{t^3}{21} \Big|_{26}^{52} = 676/3$   
Set  $676/3 = t^2/14$ ,  $t = \pm \frac{26}{3} \sqrt{21}$ , so  $t \approx 39.716$ , so during the 40th week.

$$41. \quad u = 5 + 2 \sin 3x, \quad du = 6 \cos 3x dx; \quad \int \frac{1}{6\sqrt{u}} du = \frac{1}{3} u^{1/2} + C = \frac{1}{3} \sqrt{5 + 2 \sin 3x} + C$$

$$42. \quad u = 3 + \sqrt{x}, \quad du = \frac{1}{2\sqrt{x}}dx; \quad \int 2\sqrt{u}du = \frac{4}{3}u^{3/2} + C = \frac{4}{3}(3 + \sqrt{x})^{3/2} + C$$

43.  $u = ax^3 + b, du = 3ax^2dx; \int \frac{1}{3au^2}du = -\frac{1}{3au} + C = -\frac{1}{3a^2x^3 + 3ab} + C$

44.  $u = ax^2, du = 2axdx; \frac{1}{2a} \int \sec^2 u du = \frac{1}{2a} \tan u + C = \frac{1}{2a} \tan(ax^2) + C$

45.  $\left( -\frac{1}{3u^3} - \frac{3}{u} + \frac{1}{4u^4} \right) \Big|_{-2}^{-1} = 389/192$       46.  $\frac{1}{3\pi} \sin^3 \pi x \Big|_0^1 = 0$

47. With  $b = 1.618034$ , area  $= \int_0^b (x + x^2 - x^3)dx = 1.007514$ .


48. (a)  $f(x) = \frac{1}{3}x^2 \sin 3x - \frac{2}{27} \sin 3x + \frac{2}{9}x \cos 3x - 0.251607$

(b)  $f(x) = \sqrt{4+x^2} + \frac{4}{\sqrt{4+x^2}} - 6$

49. (a) Solve  $\frac{1}{4}k^4 - k - k^2 + \frac{7}{4} = 0$  to get  $k = 2.073948$ .

(b) Solve  $-\frac{1}{2} \cos 2k + \frac{1}{3}k^3 + \frac{1}{2} = 3$  to get  $k = 1.837992$ .

50.  $F(x) = \int_{-1}^x \frac{t}{\sqrt{2+t^3}} dt, F'(x) = \frac{x}{\sqrt{2+x^3}}$ , so  $F$  is increasing on  $[1, 3]$ ;  $F_{\max} = F(3) \approx 1.152082854$   
and  $F_{\min} = F(1) \approx -0.07649493141$



52.  $\lim_{n \rightarrow +\infty} \sum_{k=1}^n \left[ \frac{25(k-1)}{n} - \frac{25(k-1)^2}{n^2} \right] \frac{5}{n} = \frac{125}{6}$

## CHAPTER 5 HORIZON MODULE

1.  $v_x(0) = 35 \cos \alpha$ , so from Equation (1),  $x(t) = (35 \cos \alpha)t$ ;  $v_y(0) = 35 \sin \alpha$ , so from Equation (2),  $y(t) = (35 \sin \alpha)t - 4.9t^2$ .


2. (a)  $v_x(t) = \frac{dx(t)}{dt} = 35 \cos \alpha, v_y(t) = \frac{dy(t)}{dt} = 35 \sin \alpha - 9.8t$

(b)  $v_y(t) = 35 \sin \alpha - 9.8t, v_y(t) = 0$  when  $t = 35 \sin \alpha / 9.8$ ;  
 $y = v_y(0)t - 4.9t^2 = (35 \sin \alpha)(35 \sin \alpha) / 9.8 - 4.9((35 \sin \alpha) / 9.8)^2 = 62.5 \sin^2 \alpha$ , so  
 $y_{\max} = 62.5 \sin^2 \alpha$ .

3.  $t = x/(35 \cos \alpha)$  so  $y = (35 \sin \alpha)(x/(35 \cos \alpha)) - 4.9(x/(35 \cos \alpha))^2 = (\tan \alpha)x - \frac{0.004}{\cos^2 \alpha}x^2$ ;  
the trajectory is a parabola because  $y$  is a quadratic function of  $x$ .

4.

$15^\circ$	$25^\circ$	$35^\circ$	$45^\circ$	$55^\circ$	$65^\circ$	$75^\circ$	$85^\circ$
no	yes	no	no	no	yes	no	no



5.  $y(t) = (35 \sin \alpha) t - 4.9t^2 = 0$  when  $t = 35 \sin \alpha / 4.9$ , at which time  
 $x = (35 \cos \alpha)(35 \sin \alpha / 4.9) = 125 \sin 2\alpha$ ; this is the maximum value of  $x$ , so  $R = 125 \sin 2\alpha$  m.
6. (a)  $R = 95$  when  $\sin 2\alpha = 95/125 = 0.76$ ,  $\alpha = 0.4316565575, 1.139139769$  rad  $\approx 24.73^\circ, 65.27^\circ$ .  
(b)  $y(t) < 50$  is required; but  $y(1.139) \approx 51.56$  m, so his height would be 56.56 m.
7.  $0.4019 < \alpha < 0.4636$  (radians), or  $23.03^\circ < \alpha < 26.57^\circ$