



The University of New Mexico

Introduction to Computer Graphics with WebGL

- Ed Angel
- Professor Emeritus of Computer Science
- Founding Director, Arts, Research, Technology and Science Laboratory
- University of New Mexico



The University of New Mexico

Geometry

- Ed Angel
- Professor Emeritus of Computer Science
- University of New Mexico



Objectives

- Introduce the elements of geometry
 - Scalars
 - Vectors
 - Points
- Develop mathematical operations among them in a coordinate-free manner
- Define basic primitives
 - Line segments
 - Polygons



Basic Elements

- Geometry is the study of the relationships among objects in an n -dimensional space

In computer graphics, we are interested in objects that exist in three dimensions

- Want a minimum set of primitives from which we can build more sophisticated objects
- We will need three basic elements

Scalars

Vectors

Points



Coordinate-Free Geometry

- When we learned simple geometry, most of us started with a Cartesian approach

Points were at locations in space $\mathbf{p}=(x,y,z)$

We derived results by algebraic manipulations involving these coordinates

- This approach was nonphysical

Physically, points exist regardless of the location of an arbitrary coordinate system

Most geometric results are independent of the coordinate system

Example Euclidean geometry: two triangles are identical if two corresponding sides and the angle between them are identical



Scalars

-
- Need three basic elements in geometry
Scalars, Vectors, Points
 - Scalars can be defined as members of sets which can be combined by two operations (addition and multiplication) obeying some fundamental axioms (associativity, commutativity, inverses)
 - Examples include the real and complex number systems under the ordinary rules with which we are familiar
 - Scalars alone have no geometric properties



Vectors

- Physical definition: a vector is a quantity with two attributes

Direction

Magnitude


- Examples include

Force

Velocity

Directed line segments


- Most important example for graphics
- Can map to other types





Vector Operations

- Every vector has a symmetric vector
Same magnitude but points in opposite direction
- Every vector can be multiplied by a scalar
- There is a zero vector
Zero magnitude, undefined orientation
- The sum of any two vectors is a vector
Use head-to-tail axiom





Linear Vector Spaces

- Mathematical system for manipulating vectors
- Operations

Scalar-vector multiplication $u = \alpha v$

Vector-vector addition: $w = u + v$

- Expressions such as


$$v = u + 2w - 3r$$

Make sense in a vector space



Vectors Lack Position

- These vectors are identical
Same length and magnitude



- Vectors spaces insufficient for geometry
Need points




Points

- Location in space
- Operations allowed between points and vectors

Point-point subtraction yields a vector

Equivalent to point-vector addition



$$v = P - Q$$

$$P = v + Q$$



Affine Spaces

- Point + a vector space
- Operations
 - Vector-vector addition
 - Scalar-vector multiplication
 - Point-vector addition
 - Scalar-scalar operations
- For any point define
 - $1 \bullet P = P$
 - $0 \bullet P = \mathbf{0}$ (zero vector)




Lines

- Consider all points of the form

$$P(\alpha) = P_0 + \alpha \mathbf{d}$$

Set of all points that pass through P_0 in the direction of the vector \mathbf{d}





Parametric Form

- This form is known as the parametric form of the line

More robust and general than other forms

Extends to curves and surfaces

- Two-dimensional forms

Explicit: $y = mx + h$

Implicit: $ax + by + c = 0$

Parametric:

$$x(\alpha) = \alpha x_0 + (1-\alpha)x_1$$

$$y(\alpha) = \alpha y_0 + (1-\alpha)y_1$$




Rays and Line Segments

- If $\alpha \geq 0$, then $P(\alpha)$ is the *ray* leaving P_0 in the direction \mathbf{d}

If we use two points to define \mathbf{v} , then

$$\begin{aligned} P(\alpha) &= Q + \alpha (R - Q) = Q + \alpha \mathbf{v} \\ &= \alpha R + (1 - \alpha)Q \end{aligned}$$


For $0 \leq \alpha \leq 1$ we get all the points on the *line segment* joining R and Q






Convexity

- An object is *convex* iff for any two points in the object all points on the line segment between these points are also in the object



convex



not convex



Affine Sums

- Consider the “sum”

$$P = \alpha_1 P_1 + \alpha_2 P_2 + \dots + \alpha_n P_n$$

Can show by induction that this sum makes sense iff

$$\alpha_1 + \alpha_2 + \dots + \alpha_n = 1$$


in which case we have the *affine sum* of the points P_1, P_2, \dots, P_n

- If, in addition, $\alpha_i \geq 0$, we have the *convex hull* of P_1, P_2, \dots, P_n



Convex Hull

- Smallest convex object containing P_1, P_2, \dots, P_n
- Formed by “shrink wrapping” points






Curves and Surfaces

- Curves are one parameter entities of the form $P(\alpha)$ where the function is nonlinear
- Surfaces are formed from two-parameter functions $P(\alpha, \beta)$

Linear functions give planes and polygons



$P(\alpha)$




$P(\alpha, \beta)$




Planes

- A plane can be defined by a point and two vectors or by three points




$$P(\alpha, \beta) = R + \alpha u + \beta v$$



$$P(\alpha, \beta) = R + \alpha(Q - R) + \beta(P - R)$$



Triangles



for $0 \leq \alpha, \beta \leq 1$, we get all points in triangle



Barycentric Coordinates

Triangle is convex so any point inside can be represented as an affine sum

$$P(\alpha_1, \alpha_2, \alpha_3) = \alpha_1 P + \alpha_2 Q + \alpha_3 R$$

where

$$\alpha_1 + \alpha_2 + \alpha_3 = 1$$

$$\alpha_i \geq 0$$

The representation is called the **barycentric coordinate** representation of P



Normals

- In three dimensional spaces, every plane has a vector n perpendicular or orthogonal to it called the **normal vector**
- From the two-point vector form $P(\alpha, \beta) = P + \alpha u + \beta v$, we know we can use the cross product to find $n = u \times v$ and the equivalent form

$$(P(\alpha, \beta) - P) \cdot n = 0$$

